arXiv:1403.5674v1 [math.AP] 22 Mar 2014

CONVERGENCE OF THE REGULARIZED SHORT PULSE EQUATION TO THE SHORT PULSE ONE

GIUSEPPE MARIA COCLITE AND LORENZO DI RUVO

Abstract. We consider the regularized short-pulse equation, which contains nonlinear dispersive effects. We prove that as the diffusion parameter tends to zero, the solutions of the dispersive equation converge to discontinuous weak solutions of the short-pulse one. The proof relies on deriving suitable a priori estimates together with an application of the compensated compactness method in the L^p setting.

1. Introduction

The nonlinear evolution equation

$$\partial_x \left(\partial_t u - \frac{1}{6} \partial_x (u^3) - \beta \partial_{xxx}^3 u \right) = \gamma u, \quad \gamma > 0, \tag{1.1}$$

known as the regularized short pulse equation, was derived by Costanzino, Manukian and Jones [9] in the context of the nonlinear Maxwell equations with high-frequency dispersion. Mathematical properties of the regularized short pulse equation (1.1) were studied recently in many details, including the local and global well-posedness in energy space [9, 16], and stability of solitary waves [9]. We rewrite (1.1) in the following way

$$\begin{cases} \partial_t u - \frac{1}{6} \partial_x (u^3) - \beta \partial_{xxx}^3 u = \gamma \displaystyle\int_0^x u(t,y) dy, & t > 0, \ x \in \mathbb{R}, \\ u(0,x) = u_0(x), & x \in \mathbb{R}, \end{cases} \tag{1.2}$$

or equivalently,

$$\begin{cases} \partial_t u - \frac{1}{6} \partial_x (u^3) - \beta \partial_{xxx}^3 u = \gamma P, & t > 0, \ x \in \mathbb{R}, \\ \partial_x P = u, & t > 0, \ x \in \mathbb{R}, \\ P(t,0) = 0, & t > 0, \\ u(0,x) = u_0(x), & x \in \mathbb{R}. \end{cases} \tag{1.3}$$

We are interested in the no high frequency limit, i.e., we send $\beta \to 0$ in (1.1). In this way, we pass from (1.1) to the equation

$$\begin{cases} \partial_x \left(\partial_t u - \frac{1}{6} \partial_x (u^3) \right) = \gamma u, & t > 0, \ x \in \mathbb{R}, \\ u(0,x) = u_0(x), & x \in \mathbb{R}. \end{cases} \tag{1.4}$$

The equation (1.4) is known as the short pulse equation, and was introduced recently by Schäfer and Wayne [20] as a model equation describing the propagation of ultra-short light pulses in silica optical fibers. It provides also an approximation of nonlinear wave packets in dispersive media in the limit of few cycles on the ultra-short pulse scale. Numerical

Date: March 25, 2014.

2000 *Mathematics Subject Classification.* 35G25, 35L65, 35L05.

Key words and phrases. Singular limit, compensated compactness, short pulse equation, entropy condition.

The authors are members of the Gruppo Nazionale per l'Analisi Matematica, la Probabilità e le loro Applicazioni (GNAMPA) of the Istituto Nazionale di Alta Matematica (INdAM).

simulations [2] show that the short pulse equation approximation to Maxwell's equations in the case when the pulse spectrum is not narrowly localized around the carrier frequency is better than the one obtained from the nonlinear Schrödinger equation, which models the evolution of slowly varying wave trains. Such ultra-short plays a key role in the development of future technologies of ultra-fast optical transmission of informations.

In [1] the author studied a new hierarchy of equations containing the short pulse equation (1.4) and the elastic beam equation, which describes nonlinear transverse oscillations of elastic beams under tension. He showed that the hierarchy of equations is integrable. He obtained the two compatible Hamiltonian structures and constructed an infinite series of both local and nonlocal conserved charges. Moreover, he gave the Lax description for both systems. The integrability and the existence of solitary wave solutions have been studied in [18, 19]. Well-posedness and wave breaking for the short pulse equation have been studied in [20] and [12], respectively.

We remark here that (1.1) and (1.4) look similar to the Ostrovsky and Ovstrosky-Hunter equations

$$\partial_x\left(\partial_t u+\frac{1}{2}\partial_x(u^2)-\beta\partial^3_{xxx}u\right)=\gamma u,\qquad \partial_x\left(\partial_t u+\frac{1}{2}\partial_x(u^2)\right)=\gamma u, \tag{1.5}$$

which was derived by Ostrovsky [15] and Hunter and Tan [11] as a model for internal solitary waves in the ocean with rotation effects of strength γ. The deep difference is in the power and the sing of the non linear term. The well-posedness on the context of discontinuous solutions of the Ovstrosky-Hunter equation has been studied in [3, 4, 7, 10]. In [6] we proved the convergence of the solutions of the Ostrovsky equation to the discontinuous ones of the Ostrovsky-Hunter equation.

Integrating (1.4) with respect to x we gain the integro-differential formulation of (1.4) (see [18])

$$\partial_t u-\partial_x\left(\frac{u^3}{6}\right)=\gamma\int_0^x u(t,y)dy,$$

that is equivalent to

$$\partial_t u-\partial_x\left(\frac{u^3}{6}\right)=\gamma P,\qquad \partial_x P=u,\quad P(\cdot,0)=0,\quad u(0,\cdot)=u_0. \tag{1.6}$$

On the initial datum, we assume that

$$u_0\in L^2(\mathbb{R})\cap L^6(\mathbb{R}),\quad \int_{\mathbb{R}}u_0(x)dx=0, \tag{1.7}$$

and on the function

$$P_0(x)=\int_{-\infty}^x u_0(y)dy,\quad x\in\mathbb{R}, \tag{1.8}$$

we assume that

$$\begin{aligned}\|P_0\|^2_{L^2(\mathbb{R})}&=\int_{\mathbb{R}}\left(\int_{-\infty}^x u_0(y)dy\right)^2dx<\infty,\\ \int_{\mathbb{R}}P_0(x)dx&=\int_{\mathbb{R}}\left(\int_{-\infty}^x u_0(y)dy\right)dx=0.\end{aligned} \tag{1.9}$$

Definition 1.1. *We say that $u\in L^\infty((0,T)\times\mathbb{R})$, $T>0$, is an entropy solution of the initial value problem* (1.4) *if*

i) u is a distributional solution of (1.6)*;*

ii) for every convex function $\eta \in C^2(\mathbb{R})$ *the entropy inequality*

$$\partial_t \eta(u) + \partial_x q(u) - \gamma \eta'(u) P \leq 0, \qquad q(u) = -\int^u \frac{\xi^2}{2} \eta'(\xi)\, d\xi \tag{1.10}$$

holds in the sense of distributions in $(0,\infty)\times\mathbb{R}$.

In [5], it is proved that (1.4) has an unique entropy solution in the sense of Definition 1.1 that depends Lipschitz continuously on the initial condition.

We study the dispersion-diffusion limit for (1.1). Therefore, we fix two small numbers $0<\varepsilon$, $\beta<1$ and consider the following third order problem

$$\begin{cases} \partial_t u_{\varepsilon,\beta} - \dfrac{1}{6}\partial_x u_{\varepsilon,\beta}^3 - \beta \partial_{xxx}^3 u_{\varepsilon,\beta} = \gamma P_{\varepsilon,\beta} + \varepsilon \partial_{xx}^2 u_{\varepsilon,\beta}, & t>0,\ x\in\mathbb{R}, \\ -\varepsilon \partial_{xx}^2 P_{\varepsilon,\beta} + \partial_x P_{\varepsilon,\beta} = u_{\varepsilon,\beta}, & t>0,\ x\in\mathbb{R}, \\ P_{\varepsilon,\beta}(t,0) = 0, & t>0, \\ u_{\varepsilon,\beta}(0,x) = u_{\varepsilon,\beta,0}(x), & x\in\mathbb{R}, \end{cases} \tag{1.11}$$

where $u_{\varepsilon,\beta,0}$ is a C^∞ approximation of u_0 such that

$$\begin{aligned} &\|u_{\varepsilon,\beta,0}\|_{L^2(\mathbb{R})}^2 + \|u_{\varepsilon,\beta,0}\|_{L^6(\mathbb{R})}^6 + (\beta+\varepsilon^2)\,\|\partial_x u_{\varepsilon,\beta,0}\|_{L^2(\mathbb{R})}^2 \leq C_0, \quad \varepsilon,\beta>0, \\ &\int_{\mathbb{R}} u_{\varepsilon,\beta,0}(x)dx = 0, \quad \varepsilon,\beta>0, \\ &\int_{\mathbb{R}} P_{0,\varepsilon,\beta}(x)dx = 0, \quad \varepsilon,\beta>0, \\ &\|P_{\varepsilon,\beta,0}\|_{L^2(\mathbb{R})}^2 + \varepsilon^2\,\|\partial_x P_{\varepsilon,\beta,0}\|_{L^2(\mathbb{R})}^2 \leq C_0, \quad \varepsilon,\beta>0, \end{aligned} \tag{1.12}$$

where C_0 is a constant independent on ε and β.

The main result of this paper is the following theorem.

Theorem 1.1. *Assume that* (1.7), (1.8), (1.9), *and* (1.12) *hold. If*

$$\beta = \mathcal{O}(\varepsilon^2), \tag{1.13}$$

then, there exist two sequences $\{\varepsilon_n\}_{n\in\mathbb{N}}$, $\{\beta_n\}_{n\in\mathbb{N}}$, *with* $\varepsilon_n, \beta_n \to 0$, *and two limit functions*

$$\begin{aligned} &u \in L^\infty(0,T;L^2(\mathbb{R})\cap L^6(\mathbb{R})), \quad T>0, \\ &P \in L^\infty((0,T)\times\mathbb{R}) \cap L^2((0,T)\times\mathbb{R}), \quad T>0, \end{aligned}$$

such that

i) u *is a distributional solution of* (1.4),
ii) $u_{\varepsilon_n,\beta_n} \to u$ *strongly in* $L^p_{loc}((0,\infty)\times\mathbb{R})$, *for each* $1\leq p<6$,
iii) $P_{\varepsilon_n,\beta_n} \to P$ *strongly in* $L^\infty((0,T)\times\mathbb{R})\cap L^2((0,T)\times\mathbb{R})$, $T>0$.

Moreover, if

$$\beta = o(\varepsilon^2), \tag{1.14}$$

then,

iv) u *is the unique entropy solution of* (1.4),
v) $P_{\varepsilon_n,\beta_n} \to P$ *strongly in* $L^p_{loc}(0,\infty;W^{1,\infty}(\mathbb{R})\cap H^1(\mathbb{R}))$, *for each* $1\leq p<6$.

In particular, we have

$$\int_{\mathbb{R}} u(t,x)dx = 0, \quad t>0. \tag{1.15}$$

The paper is organized in two sections. In Section 2, we prove some a priori estimates, while in Section 3 we prove Theorem 1.1.

2. A priori Estimates

This section is devoted to some a priori estimates on $u_{\varepsilon,\beta}$. We denote with C_0 the constants which depend only on the initial data, and with $C(T)$, the constants which depend also on T.

Arguing as [5, Lemmas 3.1, 3.2, 3.4, 3.5], we obtain the following result

Lemma 2.1. *For each $t \in (0,\infty)$,*

$$P_{\varepsilon,\beta}(t,-\infty) = \partial_x P_{\varepsilon,\beta}(t,-\infty) = P_{\varepsilon,\beta}(t,\infty) = \partial_x P_{\varepsilon,\beta}(t,\infty) = 0, \tag{2.1}$$

$$\varepsilon^2 \left\|\partial_{xx}^2 P_{\varepsilon,\beta}(t,\cdot)\right\|_{L^2(\mathbb{R})}^2 + \left\|\partial_x P_{\varepsilon,\beta}(t,\cdot)\right\|_{L^2(0,\infty)}^2 = \left\|u_{\varepsilon,\beta}(t,\cdot)\right\|_{L^2(0,\infty)}^2, \tag{2.2}$$

$$\int_{\mathbb{R}} u_{\varepsilon,\beta}(t,x)dx = 0, \tag{2.3}$$

$$\sqrt{\varepsilon}\left\|\partial_x P_{\varepsilon,\beta}(t,\cdot)\right\|_{L^\infty(\mathbb{R})} \le \left\|u_{\varepsilon,\beta}(t,\cdot)\right\|_{L^2(\mathbb{R})}, \tag{2.4}$$

$$\int_{\mathbb{R}} u_{\varepsilon,\beta}(t,x) P_{\varepsilon,\beta}(t,x)dx \le \left\|u_{\varepsilon,\beta}(t,\cdot)\right\|_{L^2(\mathbb{R})}^2. \tag{2.5}$$

Moreover, the inequality holds

$$\left\|u_{\varepsilon,\beta}(t,\cdot)\right\|_{L^2(\mathbb{R})}^2 + 2\varepsilon e^{2\gamma t}\int_0^t e^{-2\gamma s}\left\|\partial_x u_{\varepsilon,\beta}(s,\cdot)\right\|_{L^2(\mathbb{R})}^2 ds \le e^{2\gamma t} C_0. \tag{2.6}$$

In particular, we have

$$\begin{aligned} \varepsilon\left\|\partial_{xx}^2 P_{\varepsilon,\beta}(t,\cdot)\right\|_{L^2(\mathbb{R})}, \left\|\partial_x P_{\varepsilon,\beta}(t,\cdot)\right\|_{L^2(\mathbb{R})} &\le e^{\gamma t} C_0, \\ \sqrt{\varepsilon}\left\|\partial_x P_{\varepsilon,\beta}(t,\cdot)\right\|_{L^\infty(\mathbb{R})} &\le e^{\gamma t} C_0, \\ \sqrt{\varepsilon}\left|\partial_x P_{\varepsilon,\beta}(t,0)\right| &\le e^{\gamma t} C_0. \end{aligned} \tag{2.7}$$

Arguing as [6, Lemma 2.3], we have can prove the following lemma.

Lemma 2.2. *For each $t \ge 0$, we have that*

$$\int_0^{-\infty} P_{\varepsilon,\beta}(t,x)dx = a_{\varepsilon,\beta}(t), \tag{2.8}$$

$$\int_0^{\infty} P_{\varepsilon,\beta}(t,x)dx = a_{\varepsilon,\beta}(t), \tag{2.9}$$

where

$$a_{\varepsilon,\beta}(t) = \frac{1}{\gamma}\left(\varepsilon\partial_{tx}^2 P_{\varepsilon,\beta}(t,0) + \frac{1}{6}u_{\varepsilon,\beta}^3(t,0) + \beta\partial_{xx}^2 u_{\varepsilon,\beta}(t,0) + \varepsilon\partial_x u_{\varepsilon,\beta}(t,0)\right).$$

Moreover,

$$\int_{\mathbb{R}} P_{\varepsilon,\beta}(t,x)dx = 0, \quad t \ge 0. \tag{2.10}$$

Lemma 2.2 says that $P_{\varepsilon,\beta}(t,\cdot)$ is integrable at $\pm\infty$. Therefore, for each $t \ge 0$, we can consider the following function

$$F_{\varepsilon,\beta}(t,x) = \int_{-\infty}^x P_\varepsilon(t,y)dy. \tag{2.11}$$

Thus, integrating the first equation in (1.11) on $(-\infty,x)$, thanks to (2.11), we obtain that

$$\int_{-\infty}^x \partial_t u_{\varepsilon,\beta}dy - \frac{1}{6}u_{\varepsilon,\beta}^3 - \beta\partial_{xx}^2 u_{\varepsilon,\beta} - \varepsilon\partial_x u_{\varepsilon,\beta} = \gamma F_{\varepsilon,\beta}. \tag{2.12}$$

Thanks to (2.1), integrating on $\mathbb{R}$ the second equation in (1.11), we have

$$P_{\varepsilon,\beta}(t,x)=\int_{-\infty}^{x}u_{\varepsilon,\beta}(t,y)dy+\varepsilon\partial_x P_\varepsilon(t,x). \tag{2.13}$$

Differentiating (2.13) with respect to t, we get

$$\partial_t P_{\varepsilon,\beta}(t,x)=\int_{-\infty}^{x}\partial_t u_{\varepsilon,\beta}(t,y)dy+\varepsilon\partial_{tx}^2 P_{\varepsilon,\beta}(t,x). \tag{2.14}$$

Therefore, (2.12) and (2.14) give

$$\partial_t P_{\varepsilon,\beta}-\varepsilon\partial_{tx}^2 P_{\varepsilon,\beta}-\frac{1}{6}u_{\varepsilon,\beta}^3-\beta\partial_{xx}^2 u_{\varepsilon,\beta}-\varepsilon\partial_x u_{\varepsilon,\beta}=\gamma F_{\varepsilon,\beta}. \tag{2.15}$$

Lemma 2.3. *Fixed $T>0$. There exists a function $C(T)>0$, independent on ε and β, such that*

$$\|u_{\varepsilon,\beta}\|_{L^\infty((0,T)\times\mathbb{R})}\le C(T)\beta^{-\frac{1}{2}}. \tag{2.16}$$

Moreover, for every $0\le t\le T$,

$$\begin{aligned}\beta\left\|\partial_x u_{\varepsilon,\beta}(t,\cdot)\right\|_{L^2(\mathbb{R})}^2+\gamma\left\|P_{\varepsilon,\beta}(t,\cdot)\right\|_{L^2(\mathbb{R})}^2+\varepsilon^2\gamma\left\|\partial_x P_{\varepsilon,\beta}(t,\cdot)\right\|_{L^2(\mathbb{R})}^2\\ +\beta\varepsilon e^{2\gamma t}\int_0^t e^{-2\gamma s}\left\|\partial_{xx}^2 u_{\varepsilon,\beta}(s,\cdot)\right\|_{L^2(\mathbb{R})}^2 ds\le C(T)\beta^{-2}.\end{aligned} \tag{2.17}$$

Proof. Let $0\le t\le T$. Multiplying (1.11) by $-\beta\partial_{xx}^2 u_{\varepsilon,\beta}-\frac{1}{6}u_{\varepsilon,\beta}^3$, we have

$$\begin{aligned}&\left(-\beta\partial_{xx}^2 u_{\varepsilon,\beta}-\frac{1}{6}u_{\varepsilon,\beta}^3\right)\partial_t u_{\varepsilon,\beta}-\frac{1}{6}\left(-\beta\partial_{xx}^2 u_{\varepsilon,\beta}-\frac{1}{6}u_{\varepsilon,\beta}^3\right)\partial_x u_{\varepsilon,\beta}^3\\ &\qquad-\beta\left(-\beta\partial_{xx}^2 u_{\varepsilon,\beta}-\frac{1}{6}u_{\varepsilon,\beta}^3\right)\partial_{xxx}^3 u_{\varepsilon,\beta}\\ &=\gamma\left(-\beta\partial_{xx}^2 u_{\varepsilon,\beta}-\frac{1}{6}u_{\varepsilon,\beta}^3\right)P_{\varepsilon,\beta}\\ &\qquad+\varepsilon\left(-\beta\partial_{xx}^2 u_{\varepsilon,\beta}-\frac{1}{6}u_{\varepsilon,\beta}^3\right)\partial_{xx}^2 u_{\varepsilon,\beta}.\end{aligned} \tag{2.18}$$

Observe that

$$\begin{aligned}\int_{\mathbb{R}}\left(-\beta\partial_{xx}^2 u_{\varepsilon,\beta}-\frac{1}{6}u_{\varepsilon,\beta}^3\right)\partial_t u_{\varepsilon,\beta}dx&=\frac{d}{dt}\left(\frac{\beta}{2}\left\|\partial_x u_{\varepsilon,\beta}(t,\cdot)\right\|_{L^2(\mathbb{R})}^2-\frac{1}{24}\int_{\mathbb{R}}u_{\varepsilon,\beta}^4dx\right),\\ -\frac{1}{6}\int_{\mathbb{R}}\left(-\beta\partial_{xx}^2 u_{\varepsilon,\beta}-\frac{1}{6}u_{\varepsilon,\beta}^3\right)\partial_x u_{\varepsilon,\beta}^3dx&=\frac{\beta}{2}\int_{\mathbb{R}}u_{\varepsilon,\beta}^2\partial_x u_{\varepsilon,\beta}\partial_{xx}^2 u_{\varepsilon,\beta}dx,\\ -\beta\int_{\mathbb{R}}\left(-\beta\partial_{xx}^2 u_{\varepsilon,\beta}-\frac{1}{6}u_{\varepsilon,\beta}^3\right)\partial_{xxx}^3 u_{\varepsilon,\beta}dx&=-\frac{\beta}{2}\int_{\mathbb{R}}u_{\varepsilon,\beta}^2\partial_x u_{\varepsilon,\beta}\partial_{xx}^2 u_{\varepsilon,\beta}dx,\\ \varepsilon\int_{\mathbb{R}}\left(-\beta\partial_{xx}^2 u_{\varepsilon,\beta}-\frac{1}{6}u_{\varepsilon,\beta}^3\right)\partial_{xx}^2 u_{\varepsilon,\beta}dx&=-\varepsilon\beta\left\|\partial_{xx}^2 u_{\varepsilon,\beta}(t,\cdot)\right\|_{L^2(\mathbb{R})}^2+\frac{\varepsilon}{2}\int_{\mathbb{R}}u_{\varepsilon,\beta}^2(\partial_x u_{\varepsilon,\beta})^2dx.\end{aligned}$$

Moreover, for (2.1),

$$-\gamma\beta\int_{\mathbb{R}}\partial_{xx}^2 u_{\varepsilon,\beta}P_{\varepsilon,\beta}dx=\gamma\beta\int_{\mathbb{R}}\partial_x u_{\varepsilon,\beta}\partial_x P_{\varepsilon,\beta}dx.$$

Therefore, integrating (2.18) over $\mathbb{R}$,

$$\begin{aligned}&\frac{d}{dt}\left(\beta\left\|\partial_x u_{\varepsilon,\beta}(t,\cdot)\right\|^2_{L^2(\mathbb{R})}-\frac{1}{12}\int_{\mathbb{R}}u^4_{\varepsilon,\beta}dx\right)+2\beta\varepsilon\left\|\partial^2_{xx}u_{\varepsilon,\beta}(t,\cdot)\right\|^2_{L^2(\mathbb{R})}\\&\quad=2\gamma\beta\int_{\mathbb{R}}\partial_x u_{\varepsilon,\beta}\partial_x P_{\varepsilon,\beta}dx-\frac{\gamma}{3}\int_{\mathbb{R}}u^3_{\varepsilon,\beta}P_{\varepsilon,\beta}dx+\varepsilon\int_{\mathbb{R}}u^2_{\varepsilon,\beta}(\partial_x u_{\varepsilon,\beta})^2dx.\end{aligned}\tag{2.19}$$

Multiplying (2.15) by $2\gamma P_{\varepsilon,\beta}-2\varepsilon\gamma\partial_x P_{\varepsilon,\beta}$, we have

$$\begin{aligned}&(\partial_t P_{\varepsilon,\beta}-\varepsilon\partial^2_{tx}P_{\varepsilon,\beta})(2\gamma P_{\varepsilon,\beta}-2\varepsilon\gamma\partial_x P_{\varepsilon,\beta})\\&\quad=\gamma F_{\varepsilon,\beta}(2\gamma P_{\varepsilon,\beta}-2\varepsilon\gamma\partial_x P_{\varepsilon,\beta})+\frac{1}{6}u^3_{\varepsilon,\beta}(2\gamma P_{\varepsilon,\beta}-2\varepsilon\gamma\partial_x P_{\varepsilon,\beta})\\&\qquad+\beta\partial^2_{xx}u_{\varepsilon,\beta}(2\gamma P_{\varepsilon,\beta}-2\varepsilon\gamma\partial_x P_{\varepsilon,\beta})+\varepsilon\partial_x u_{\varepsilon,\beta}(2\gamma P_{\varepsilon,\beta}-2\varepsilon\gamma\partial_x P_{\varepsilon,\beta}).\end{aligned}\tag{2.20}$$

It follows from (2.1) that

$$\begin{aligned}-2\varepsilon\gamma\int_{\mathbb{R}}P_{\varepsilon,\beta}\partial^2_{tx}P_{\varepsilon,\beta}dx=&2\varepsilon\gamma\int_{\mathbb{R}}\partial_x P_{\varepsilon,\beta}\partial_t P_{\varepsilon,\beta}dx,\\2\beta\gamma\int_{\mathbb{R}}\partial^2_{xx}u_{\varepsilon,\beta}P_{\varepsilon,\beta}dx=&-2\beta\gamma\int_{\mathbb{R}}\partial_x u_{\varepsilon,\beta}\partial_x P_{\varepsilon,\beta}dx.\end{aligned}\tag{2.21}$$

Therefore, arguing as [5, Lemma 2.6], an integration on $\mathbb{R}$ of (2.20), (2.1) and (2.21) give

$$\begin{aligned}&\frac{d}{dt}\left(\gamma\left\|P_{\varepsilon,\beta}(t,\cdot)\right\|^2_{L^2(\mathbb{R})}+\varepsilon^2\gamma\left\|\partial_x P_{\varepsilon,\beta}(t,\cdot)\right\|^2_{L^2(\mathbb{R})}\right)\\&\quad=2\gamma^2\int_{\mathbb{R}}F_{\varepsilon,\beta}P_{\varepsilon,\beta}dx-2\varepsilon\gamma^2\int_{\mathbb{R}}F_{\varepsilon,\beta}\partial_x P_{\varepsilon,\beta}dx\\&\qquad+\frac{\gamma}{3}\int_{\mathbb{R}}u^3_{\varepsilon,\beta}P_{\varepsilon,\beta}dx-\frac{\varepsilon\gamma}{3}\int_{\mathbb{R}}u^3_{\varepsilon,\beta}\partial_x P_{\varepsilon,\beta}dx\\&\qquad-2\beta\gamma\int_{\mathbb{R}}\partial_x u_{\varepsilon,\beta}\partial_x P_{\varepsilon,\beta}dx-2\beta\varepsilon\int_{\mathbb{R}}\partial^2_{xx}u_{\varepsilon,\beta}P_{\varepsilon,\beta}dx\\&\qquad+2\varepsilon\gamma\int_{\mathbb{R}}\partial_x u_{\varepsilon,\beta}P_{\varepsilon,\beta}dx-2\varepsilon^2\gamma\int_{\mathbb{R}}\partial_x u_{\varepsilon,\beta}\partial_x P_{\varepsilon,\beta}dx.\end{aligned}\tag{2.22}$$

Due to (2.10) and (2.11),

$$\begin{aligned}2\gamma^2\int_{\mathbb{R}}F_{\varepsilon,\beta}P_{\varepsilon,\beta}dx=&2\gamma^2\int_{\mathbb{R}}F_{\varepsilon,\beta}\partial_x F_{\varepsilon,\beta}dx\\=&\gamma^2(F_{\varepsilon,\beta}(t,\infty))^2=\gamma^2\left(\int_{\mathbb{R}}P_{\varepsilon,\beta}(t,x)dx\right)^2=0.\end{aligned}\tag{2.23}$$

It follows from (2.22) and (2.23) that

$$\begin{aligned}&\frac{d}{dt}\left(\gamma\left\|P_{\varepsilon,\beta}(t,\cdot)\right\|^2_{L^2(\mathbb{R})}+\varepsilon^2\gamma\left\|\partial_x P_{\varepsilon,\beta}(t,\cdot)\right\|^2_{L^2(\mathbb{R})}\right)\\&\quad=-2\varepsilon\gamma^2\int_{\mathbb{R}}F_{\varepsilon,\beta}\partial_x P_{\varepsilon,\beta}dx+\frac{\gamma}{3}\int_{\mathbb{R}}u^3_{\varepsilon,\beta}P_{\varepsilon,\beta}dx\\&\qquad-\frac{\varepsilon\gamma}{3}\int_{\mathbb{R}}u^3_{\varepsilon,\beta}\partial_x P_{\varepsilon,\beta}dx-2\beta\gamma\int_{\mathbb{R}}\partial_x u_{\varepsilon,\beta}\partial_x P_{\varepsilon,\beta}dx\\&\qquad-2\beta\varepsilon\int_{\mathbb{R}}\partial_x u_{\varepsilon,\beta}\partial_x P_{\varepsilon,\beta}dx+2\varepsilon\gamma\int_{\mathbb{R}}\partial_x u_{\varepsilon,\beta}P_{\varepsilon,\beta}dx\\&\qquad-2\varepsilon^2\gamma\int_{\mathbb{R}}\partial_x u_{\varepsilon,\beta}\partial_x P_{\varepsilon,\beta}dx.\end{aligned}\tag{2.24}$$

Adding (2.19) and (2.24), we get

$$\begin{aligned}
&\frac{d}{dt}G_1(t) + 2\beta\varepsilon \left\|\partial_{xx}^2 u_{\varepsilon,\beta}(t,\cdot)\right\|_{L^2(\mathbb{R})}^2 \\
&\quad = \varepsilon \int_{\mathbb{R}} u_{\varepsilon,\beta}^2 (\partial_x u_{\varepsilon,\beta})^2 dx - 2\varepsilon\gamma^2 \int_{\mathbb{R}} F_{\varepsilon,\beta}\partial_x P_{\varepsilon,\beta} dx \\
&\quad - \frac{\varepsilon\gamma}{3}\int_{\mathbb{R}} u_{\varepsilon,\beta}^3 \partial_x P_{\varepsilon,\beta} dx - 2\varepsilon\beta \int_{\mathbb{R}} \partial_{xx}^2 u_{\varepsilon,\beta} P_{\varepsilon,\beta} dx \\
&\quad + 2\varepsilon\gamma \int_{\mathbb{R}} \partial_x u_{\varepsilon,\beta} P_{\varepsilon,\beta} dx - 2\varepsilon^2\gamma \int_{\mathbb{R}} \partial_x u_{\varepsilon,\beta}\partial_x P_{\varepsilon,\beta} dx,
\end{aligned} \tag{2.25}$$

where

$$\begin{aligned}
G_1(t) = \beta \left\|\partial_x u_{\varepsilon,\beta}(t,\cdot)\right\|_{L^2(\mathbb{R})}^2 - \frac{1}{12}\int_{\mathbb{R}} u_{\varepsilon,\beta}^4 dx \\
+ \gamma \left\|P_{\varepsilon,\beta}(t,\cdot)\right\|_{L^2(\mathbb{R})}^2 + \varepsilon^2\gamma \left\|\partial_x P_{\varepsilon,\beta}(t,\cdot)\right\|_{L^2(\mathbb{R})}^2.
\end{aligned} \tag{2.26}$$

Since $0 < \varepsilon < 1$, thanks to (2.1) and (2.11),

$$-2\varepsilon\gamma^2 \int_{\mathbb{R}} F_{\varepsilon,\beta}\partial_x P_{\varepsilon,\beta} dx = 2\varepsilon\gamma^2 \int_{\mathbb{R}} \partial_x F_{\varepsilon,\beta} P_{\varepsilon,\beta} dx \le 2\gamma^2 \left\|P_{\varepsilon,\beta}(t,\cdot)\right\|_{L^2(\mathbb{R})}^2. \tag{2.27}$$

Since $0 < \varepsilon, \beta < 1$, due to (2.1), (2.7) and the Young inequality,

$$\begin{aligned}
&-2\varepsilon\beta \int_{\mathbb{R}} \partial_{xx}^2 u_{\varepsilon,\beta} P_{\varepsilon,\beta} dx = 2\varepsilon\beta \int_{\mathbb{R}} \partial_x u_{\varepsilon,\beta}\partial_x P_{\varepsilon,\beta} dx \\
&\le 2\varepsilon\beta \left|\int_{\mathbb{R}} \partial_x u_{\varepsilon,\beta}\partial_x P_{\varepsilon,\beta} dx\right| \\
&\le 2\varepsilon \int_{\mathbb{R}} |\partial_x u_{\varepsilon,\beta}||\partial_x P_{\varepsilon}| dx \\
&\le \varepsilon \left\|\partial_x u_{\varepsilon,\beta}(t,\cdot)\right\|_{L^2(\mathbb{R})}^2 + \varepsilon \left\|P_{\varepsilon,\beta}(t,\cdot)\right\|_{L^2(\mathbb{R})}^2 \\
&\le \varepsilon \left\|\partial_x u_{\varepsilon,\beta}(t,\cdot)\right\|_{L^2(\mathbb{R})}^2 + \left\|P_{\varepsilon,\beta}(t,\cdot)\right\|_{L^2(\mathbb{R})}^2, \\
&\le \varepsilon \left\|\partial_x u_{\varepsilon,\beta}(t,\cdot)\right\|_{L^2(\mathbb{R})}^2 + C_0 e^{2\gamma t} \\
&\le \varepsilon \left\|\partial_x u_{\varepsilon,\beta}(t,\cdot)\right\|_{L^2(\mathbb{R})}^2 + C(T),
\end{aligned} \tag{2.28}$$

$$\begin{aligned}
&2\varepsilon\gamma \int_{\mathbb{R}} \partial_x u_{\varepsilon,\beta} P_{\varepsilon,\beta} dx = -2\varepsilon\gamma \int_{\mathbb{R}} u_{\varepsilon,\beta}\partial_x P_{\varepsilon,\beta} dx \\
&\le 2\varepsilon\gamma \left|\int_{\mathbb{R}} u_{\varepsilon,\beta}\partial_x P_{\varepsilon,\beta} dx\right| \\
&\le 2\varepsilon\gamma \int_{\mathbb{R}} |u_{\varepsilon,\beta}||\partial_x P_{\varepsilon,\beta}| dx \\
&\le \varepsilon\gamma \left\|u_{\varepsilon,\beta}(t,\cdot)\right\|_{L^2(\mathbb{R})}^2 + \varepsilon\gamma \left\|\partial_x P_{\varepsilon,\beta}(t,\cdot)\right\|_{L^2(\mathbb{R})}^2 \\
&\le \gamma \left\|u_{\varepsilon,\beta}(t,\cdot)\right\|_{L^2(\mathbb{R})}^2 + \gamma \left\|\partial_x P_{\varepsilon,\beta}(t,\cdot)\right\|_{L^2(\mathbb{R})}^2 \\
&\le \gamma C_0 e^{2\gamma t} + \gamma C_0 e^{2\gamma t} \le C(T).
\end{aligned}$$

For the Young inequality,

$$
\begin{aligned}
2\varepsilon^2\gamma\left|\int_{\mathbb{R}}\partial_x u_{\varepsilon,\beta}\partial_x P_{\varepsilon,\beta}dx\right| \leq & \varepsilon^2\int_{\mathbb{R}}|\partial_x u_{\varepsilon,\beta}||2\gamma\partial_x P_{\varepsilon,\beta}|dx\\
\leq & \frac{\varepsilon^2}{2}\left\|\partial_x u_{\varepsilon,\beta}(t,\cdot)\right\|^2_{L^2(\mathbb{R})}+2\varepsilon^2\gamma^2\left\|\partial_x P_{\varepsilon,\beta}(t,\cdot)\right\|^2_{L^2(\mathbb{R})}\\
\leq & \frac{\varepsilon}{2}\left\|\partial_x u_{\varepsilon,\beta}(t,\cdot)\right\|^2_{L^2(\mathbb{R})}+2\varepsilon^2\gamma^2\left\|\partial_x P_{\varepsilon,\beta}(t,\cdot)\right\|^2_{L^2(\mathbb{R})}.
\end{aligned}
\tag{2.29}
$$

It follows from (2.25), (2.26), (2.27), (2.28) and (2.29) that

$$
\begin{aligned}
&\frac{d}{dt}G_1(t)+2\beta\varepsilon\left\|\partial^2_{xx}u_{\varepsilon,\beta}(t,\cdot)\right\|^2_{L^2(\mathbb{R})}\\
&\quad\leq 2\gamma^2\left\|P_{\varepsilon,\beta}(t,\cdot)\right\|^2_{L^2(\mathbb{R})}+2\gamma^2\varepsilon^2\left\|\partial_x P_{\varepsilon,\beta}(t,\cdot)\right\|^2_{L^2(\mathbb{R})}\\
&\qquad+\varepsilon\int_{\mathbb{R}}u^2_{\varepsilon,\beta}(\partial_x u_{\varepsilon,\beta})^2dx+\frac{\varepsilon\gamma}{3}\int_{\mathbb{R}}|u_{\varepsilon,\beta}|^3||\partial_x P_{\varepsilon,\beta}|dx\\
&\qquad+\frac{3\varepsilon}{2}\left\|\partial_x u_{\varepsilon,\beta}(t,\cdot)\right\|^2_{L^2(\mathbb{R})}+C(T)\\
&\quad\leq 2\gamma\beta\left\|\partial_x u_{\varepsilon,\beta}(t,\cdot)\right\|^2_{L^2(\mathbb{R})}+2\gamma^2\left\|P_{\varepsilon,\beta}(t,\cdot)\right\|^2_{L^2(\mathbb{R})}\\
&\qquad+2\gamma^2\varepsilon^2\left\|\partial_x P_{\varepsilon,\beta}(t,\cdot)\right\|^2_{L^2(\mathbb{R})}+\varepsilon\int_{\mathbb{R}}u^2_{\varepsilon,\beta}(\partial_x u_{\varepsilon,\beta})^2dx\\
&\qquad+\frac{\varepsilon\gamma}{3}\int_{\mathbb{R}}|u_{\varepsilon,\beta}|^3||\partial_x P_{\varepsilon,\beta}|dx-\frac{\gamma}{6}\int_{\mathbb{R}}u^4_{\varepsilon,\beta}dx\\
&\qquad+\frac{\gamma}{6}\int_{\mathbb{R}}u^4_{\varepsilon,\beta}dx+\frac{3\varepsilon}{2}\left\|\partial_x u_{\varepsilon,\beta}(t,\cdot)\right\|^2_{L^2(\mathbb{R})}+C(T)\\
&\quad=2\gamma G_1(t)+\varepsilon\int_{\mathbb{R}}u^2_{\varepsilon,\beta}(\partial_x u_{\varepsilon,\beta})^2dx+\frac{\varepsilon\gamma}{3}\int_{\mathbb{R}}|u_{\varepsilon,\beta}|^3||\partial_x P_{\varepsilon,\beta}|dx\\
&\qquad+\frac{\gamma}{6}\int_{\mathbb{R}}u^4_{\varepsilon,\beta}dx+\frac{3\varepsilon}{2}\left\|\partial_x u_{\varepsilon,\beta}(t,\cdot)\right\|^2_{L^2(\mathbb{R})}+C(T),
\end{aligned}
$$

that is

$$
\begin{aligned}
&\frac{d}{dt}G_1(t)-2\gamma G_1(t)+2\beta\varepsilon\left\|\partial^2_{xx}u_{\varepsilon,\beta}(t,\cdot)\right\|^2_{L^2(\mathbb{R})}\\
&\qquad\leq\varepsilon\int_{\mathbb{R}}u^2_{\varepsilon,\beta}(\partial_x u_{\varepsilon,\beta})^2dx+\frac{\varepsilon\gamma}{3}\int_{\mathbb{R}}|u_{\varepsilon,\beta}|^3||\partial_x P_{\varepsilon,\beta}|dx\\
&\qquad\quad+\frac{\gamma}{6}\int_{\mathbb{R}}u^4_{\varepsilon,\beta}dx+\frac{3\varepsilon}{2}\left\|\partial_x u_{\varepsilon,\beta}(t,\cdot)\right\|^2_{L^2(\mathbb{R})}+C(T).
\end{aligned}
\tag{2.30}
$$

Since $0<\varepsilon<1$, due to (2.6), (2.7) and the Young inequality,

$$
\begin{aligned}
&\frac{\varepsilon\gamma}{3}\int_{\mathbb{R}}|u_{\varepsilon,\beta}|^3||\partial_x P_{\varepsilon,\beta}|dx\leq\frac{\varepsilon\gamma}{3}\int_{\mathbb{R}}u^2_{\varepsilon,\beta}(\partial_x P_{\varepsilon,\beta})^2dx+\frac{\varepsilon\gamma}{3}\int_{\mathbb{R}}u^4_{\varepsilon,\beta}dx\\
&\quad\leq\frac{\gamma}{3}\varepsilon\left\|\partial_x P_{\varepsilon,\beta}(t,\cdot)\right\|^2_{L^\infty(\mathbb{R})}\left\|u_{\varepsilon,\beta}(t,\cdot)\right\|^2_{L^2(\mathbb{R})}+\frac{\gamma}{3}\left\|u_{\varepsilon,\beta}\right\|^2_{L^\infty((0,T)\times\mathbb{R})}\left\|u_{\varepsilon,\beta}(t,\cdot)\right\|^2_{L^2(\mathbb{R})}\\
&\quad\leq\frac{\gamma}{3}C_0e^{4\gamma t}+\frac{\gamma}{3}C_0e^{2\gamma t}\left\|u_{\varepsilon,\beta}\right\|^2_{L^\infty((0,T)\times\mathbb{R})}\leq C(T)+C(T)\left\|u_{\varepsilon,\beta}\right\|^2_{L^\infty((0,T)\times\mathbb{R})}.
\end{aligned}
\tag{2.31}
$$

Thanks to (2.6),

$$
\begin{aligned}
\frac{\gamma}{6}\int_{\mathbb{R}}u^4_{\varepsilon,\beta}dx\leq&\frac{\gamma}{6}\left\|u_{\varepsilon,\beta}\right\|^2_{L^\infty((0,T)\times\mathbb{R})}\left\|u_{\varepsilon,\beta}(t,\cdot)\right\|^2_{L^2(\mathbb{R})}\\
\leq&\frac{\gamma}{6}C_0e^{2\gamma t}\left\|u_{\varepsilon,\beta}\right\|^2_{L^\infty((0,T)\times\mathbb{R})}\leq C(T)\left\|u_{\varepsilon,\beta}\right\|^2_{L^\infty((0,T)\times\mathbb{R})}.
\end{aligned}
\tag{2.32}
$$

(2.30), (2.31) and (2.32) give

$$
\begin{aligned}
&\frac{d}{dt}G_1(t) - 2\gamma G_1(t) + 2\beta\varepsilon \left\|\partial_{xx}^2 u_{\varepsilon,\beta}(t,\cdot)\right\|_{L^2(\mathbb{R})}^2 \\
&\qquad \leq C(T) + C(T)\left\|u_{\varepsilon,\beta}\right\|_{L^\infty((0,T)\times\mathbb{R})}^2 + \varepsilon\left\|u_{\varepsilon,\beta}\right\|_{L^\infty((0,T)\times\mathbb{R})}^2 \left\|\partial_x u_{\varepsilon,\beta}(t,\cdot)\right\|_{L^2(\mathbb{R})}^2 \\
&\qquad\quad + \frac{3\varepsilon}{2}\left\|\partial_x u_{\varepsilon,\beta}(t,\cdot)\right\|_{L^2(\mathbb{R})}^2 .
\end{aligned} \tag{2.33}
$$

It follows from (1.12), (2.6), (2.26), (2.33) and Gronwall's Lemma that

$$
\begin{aligned}
&\beta\left\|\partial_x u_{\varepsilon,\beta}(t,\cdot)\right\|_{L^2(\mathbb{R})}^2 + \gamma\left\|P_{\varepsilon,\beta}(t,\cdot)\right\|_{L^2(\mathbb{R})}^2 \\
&\qquad + \varepsilon^2\gamma\left\|\partial_x P_{\varepsilon,\beta}(t,\cdot)\right\|_{L^2(\mathbb{R})}^2 + 2\beta\varepsilon e^{2\gamma t}\int_0^t e^{-2C\gamma s}\left\|\partial_{xx}^2 u_{\varepsilon,\beta}(s,\cdot)\right\|_{L^2(\mathbb{R})}^2 ds \\
&\leq C_0 e^{2\gamma t} + C(T)e^{2\gamma t}\int_0^t e^{-2\gamma s}ds + e^{2\gamma t}C(T)\left\|u_{\varepsilon,\beta}\right\|_{L^\infty((0,T)\times\mathbb{R})}^2\int_0^t e^{-2\gamma s}ds \\
&\qquad + \varepsilon e^{2\gamma t}\left\|u_{\varepsilon,\beta}\right\|_{L^\infty((0,T)\times\mathbb{R})}^2\int_0^t e^{-2\gamma s}ds\left\|\partial_x u_{\varepsilon,\beta}(s,\cdot)\right\|_{L^2(\mathbb{R})}^2 ds \\
&\qquad + \frac{3\varepsilon}{2}e^{2\gamma t}\int_0^t e^{-2\gamma s}\left\|\partial_x u_{\varepsilon,\beta}(s,\cdot)\right\|_{L^2(\mathbb{R})}^2 ds + \frac{1}{12}\int_{\mathbb{R}} u_{\varepsilon,\beta}^4 dx \\
&\leq C(T) + C(T)\left\|u_{\varepsilon,\beta}\right\|_{L^\infty((0,T)\times\mathbb{R})}^2 + \frac{C_0 e^{2\gamma t}}{2}\left\|u_{\varepsilon,\beta}\right\|_{L^\infty((0,T)\times\mathbb{R})}^2 \\
&\quad 3C_0 e^{2\gamma t} + \frac{1}{12}\left\|u_{\varepsilon,\beta}\right\|_{L^\infty((0,T)\times\mathbb{R})}^2\left\|u_{\varepsilon,\beta}(t,\cdot)\right\|_{L^2(\mathbb{R})}^2 \\
&\leq C(T) + C(T)\left\|u_{\varepsilon,\beta}\right\|_{L^\infty((0,T)\times\mathbb{R})}^2 + \frac{C_0 e^{2\gamma t}}{12}\left\|u_{\varepsilon,\beta}\right\|_{L^\infty((0,T)\times\mathbb{R})}^2 \\
&\leq C(T)\left(1 + \left\|u_{\varepsilon,\beta}\right\|_{L^\infty((0,T)\times\mathbb{R})}^2\right).
\end{aligned} \tag{2.34}
$$

Following [6, 8, 13], we begin by observing that, due to (2.6), (2.34) and the Hölder inequality,

$$
\begin{aligned}
u_{\varepsilon,\beta}^2(t,x) &= 2\int_{-\infty}^x u_{\varepsilon,\beta}\partial_x u_{\varepsilon,\beta}dy \leq 2\int_{\mathbb{R}}|u_{\varepsilon,\beta}\partial_x u_{\varepsilon,\beta}|dx \\
&\leq \frac{2}{\sqrt{\beta}}\left\|u_{\varepsilon,\beta}\right\|_{L^2(\mathbb{R})}\sqrt{\beta}\left\|\partial_x u_{\varepsilon,\beta}(t,\cdot)\right\|_{L^2(\mathbb{R})} \\
&\leq \frac{2}{\sqrt{\beta}}C_0 e^{\gamma t}\sqrt{C(T)\left(1 + \left\|u_{\varepsilon,\beta}\right\|_{L^\infty((0,T)\times\mathbb{R})}^2\right)} \\
&\leq \frac{C(T)}{\sqrt{\beta}}\sqrt{\left(1 + \left\|u_{\varepsilon,\beta}\right\|_{L^\infty((0,T)\times\mathbb{R})}^2\right)},
\end{aligned}
$$

that is

$$
\left\|u_{\varepsilon,\beta}\right\|_{L^\infty((0,T)\times\mathbb{R})}^4 \leq \frac{C(T)}{\beta}\left(1 + \left\|u_{\varepsilon,\beta}\right\|_{L^\infty((0,T)\times\mathbb{R})}^2\right). \tag{2.35}
$$

Let us show that (2.35) is verified when

$$
\left\|u_{\varepsilon,\beta}\right\|_{L^\infty((0,T)\times\mathbb{R})} \leq \max\left\{1, \left(\frac{2C(T)}{\beta}\right)^{\frac{1}{2}}\right\} \leq C(T)\beta^{-\frac{1}{2}}. \tag{2.36}
$$

We write $y = \|u_{\varepsilon,\beta}\|_{L^\infty((0,T)\times\mathbb{R})}$. Therefore, it follows from (2.35) that

$$y^4 \leq \frac{C(T)}{\beta}\left(1+y^2\right). \tag{2.37}$$

If $y \leq 1$, we have (2.36).

If $y > 1$ and

$$y > \left(\frac{2C(T)}{\beta}\right)^{\frac{1}{2}},$$

that is

$$y^2 > \frac{2C(T)}{\beta}, \quad y^4 > \frac{2C(T)}{\beta}y^2. \tag{2.38}$$

It follows from (2.38) that

$$2y^4 > y^4 + y^2 > \frac{2C(T)}{\beta}(y^2+1).$$

Therefore,

$$y^4 > \frac{C(T)}{\beta}(y^2+1),$$

which is in contradiction with (2.37).

Finally, (2.17) follows from (2.16) and (2.34). □

Lemma 2.4. *Let $T > 0$. Assume (1.13) holds true. There exists a function $C(T) > 0$, independent on ε and β, such that*

$$\|P_{\varepsilon,\beta}\|_{L^\infty((0,T)\times\mathbb{R})} \leq C(T). \tag{2.39}$$

In particular, we have that

$$\|P_{\varepsilon,\beta}(t,\cdot)\|_{L^2(\mathbb{R})} \leq C(T), \tag{2.40}$$

$$\varepsilon\,\|\partial_x P_{\varepsilon,\beta}(t,\cdot)\|_{L^2(\mathbb{R})} \leq C(T), \tag{2.41}$$

$$\|u_{\varepsilon,\beta}(t,\cdot)\|_{L^6(\mathbb{R})} \leq C(T), \tag{2.42}$$

$$\varepsilon\,\|\partial_x u_{\varepsilon,\beta}(t,\cdot)\|_{L^2(\mathbb{R})} \leq C(T), \tag{2.43}$$

$$\varepsilon e^t \int_0^t\int_{\mathbb{R}} e^{-s} u_{\varepsilon,\beta}^4(s,\cdot)(\partial_x u_{\varepsilon,\beta}(s,\cdot))^2 ds dx \leq C(T), \tag{2.44}$$

$$\varepsilon^3 e^t \int_0^t e^{-s}\left\|\partial_{xx}^2 u_{\varepsilon,\beta}(s,\cdot)\right\|_{L^2(\mathbb{R})}^2 ds \leq C(T), \tag{2.45}$$

for every $0 < t < T$. Moreover,

$$\beta\left\|\partial_x u_{\varepsilon,\beta}\partial_{xx}^2 u_{\varepsilon,\beta}\right\|_{L^1((0,T)\times\mathbb{R})} \leq C(T), \tag{2.46}$$

$$\beta^2 \int_0^T \left\|\partial_{xx}^2 u_{\varepsilon,\beta}(s,\cdot)\right\|_{L^2(\mathbb{R})}^2 ds \leq C(T)\varepsilon. \tag{2.47}$$

Proof. Let $0 \leq t \leq T$. Multiplying (1.11) by $u_{\varepsilon,\beta}^5 - 3\varepsilon^2\partial_{xx}^2 u_{\varepsilon,\beta}$, we have

$$\begin{aligned}(u_{\varepsilon,\beta}^5 - 3\varepsilon^2\partial_{xx}^2 u_{\varepsilon,\beta})\partial_t u_{\varepsilon,\beta} - \frac{1}{6}(u_{\varepsilon,\beta}^5 - 3\varepsilon^2\partial_{xx}^2 u_{\varepsilon,\beta})\partial_x u_{\varepsilon,\beta}^3 \\ - (u_{\varepsilon,\beta}^5 - 3\varepsilon^2\partial_{xx}^2 u_{\varepsilon,\beta})\beta\partial_{xxx}^3 u_{\varepsilon,\beta} \\ = \gamma(u_{\varepsilon,\beta}^5 - 3\varepsilon^2\partial_{xx}^2 u_{\varepsilon,\beta})P_{\varepsilon,\beta} + \varepsilon(u_{\varepsilon,\beta}^5 - 3\varepsilon^2\partial_{xx}^2 u_{\varepsilon,\beta})\partial_{xx}^2 u_{\varepsilon,\beta}.\end{aligned} \tag{2.48}$$

Since

$$\begin{aligned}
\int_{\mathbb{R}}(u_{\varepsilon,\beta}^5-3\varepsilon^2\partial_{xx}^2u_{\varepsilon,\beta})\partial_t u_{\varepsilon,\beta}dx =&\frac{d}{dt}\left(\frac{1}{6}\left\|u_{\varepsilon,\beta}(t,\cdot)\right\|_{L^6(\mathbb{R})}^6+\frac{3\varepsilon^2}{2}\left\|\partial_x u_{\varepsilon,\beta}(t,\cdot)\right\|_{L^2(\mathbb{R})}^2\right),\\
-\frac{1}{6}\int_{\mathbb{R}}(u_{\varepsilon,\beta}^5-3\varepsilon^2\partial_{xx}^2u_{\varepsilon,\beta})\partial_x u_{\varepsilon,\beta}^3dx =&\frac{3\varepsilon^2}{2}\int_{\mathbb{R}}u_{\varepsilon,\beta}^2\partial_x u_{\varepsilon,\beta}\partial_{xx}^2u_{\varepsilon,\beta}dx,\\
-\beta\int_{\mathbb{R}}(u_{\varepsilon,\beta}^5-3\varepsilon^2\partial_{xx}^2u_{\varepsilon,\beta})\partial_{xxx}^3u_{\varepsilon,\beta}dx =&5\beta\int_{\mathbb{R}}u_{\varepsilon,\beta}^4\partial_x u_{\varepsilon,\beta}\partial_{xx}^2u_{\varepsilon,\beta}dx\\
=&-10\beta\int_{\mathbb{R}}u_{\varepsilon,\beta}^3(\partial_x u_{\varepsilon,\beta})^2dx,\\
\varepsilon\int_{\mathbb{R}}(u_{\varepsilon,\beta}^5-3\varepsilon^2\partial_{xx}^2u_{\varepsilon,\beta})\partial_{xx}^2u_{\varepsilon,\beta}dx =&-5\varepsilon\int_{\mathbb{R}}u_{\varepsilon,\beta}^4(\partial_x u_{\varepsilon,\beta})^2dx-3\varepsilon^3\left\|\partial_{xx}^2u_{\varepsilon,\beta}(t,\cdot)\right\|_{L^2(\mathbb{R})}^2,
\end{aligned}$$

integrating (2.48) over $\mathbb{R}$,

$$\begin{aligned}
&\frac{d}{dt}\left(\frac{1}{6}\left\|u_{\varepsilon,\beta}(t,\cdot)\right\|_{L^6(\mathbb{R})}^6+\frac{3\varepsilon^2}{2}\left\|\partial_x u_{\varepsilon,\beta}(t,\cdot)\right\|_{L^2(\mathbb{R})}^2\right)\\
&\quad+5\varepsilon\int_{\mathbb{R}}u_{\varepsilon,\beta}^4(\partial_x u_{\varepsilon,\beta})^2dx+3\varepsilon^3\left\|\partial_{xx}^2u_{\varepsilon,\beta}(t\cdot)\right\|_{L^2(\mathbb{R})}^2\\
&=\gamma\int_{\mathbb{R}}u_{\varepsilon,\beta}^5P_{\varepsilon,\beta}dx-3\gamma\varepsilon^2\int_{\mathbb{R}}\partial_{xx}^2u_{\varepsilon,\beta}P_{\varepsilon,\beta}dx\\
&\quad-\frac{3\varepsilon^2}{2}\int_{\mathbb{R}}u_{\varepsilon,\beta}^2\partial_x u_{\varepsilon,\beta}\partial_{xx}^2u_{\varepsilon,\beta}dx-10\beta\int_{\mathbb{R}}u_{\varepsilon,\beta}^3(\partial_x u_{\varepsilon,\beta})^2dx.
\end{aligned}\tag{2.49}$$

Due to (2.6) and the Young inequality,

$$\begin{aligned}
\gamma\left|\int_{\mathbb{R}}u_{\varepsilon,\beta}^5P_{\varepsilon,\beta}dx\right|&\le\int_{\mathbb{R}}\left|\sqrt{6}\gamma u_{\varepsilon,\beta}^2P_{\varepsilon,\beta}\right|\left|\frac{u_{\varepsilon,\beta}^3}{\sqrt{6}}\right|dx\\
&\le3\gamma^2\int_{\mathbb{R}}P_{\varepsilon,\beta}^2u_{\varepsilon,\beta}^4dx+\frac{1}{12}\left\|u_{\varepsilon,\beta}(t,\cdot)\right\|_{L^6(\mathbb{R})}^6\\
&\le\int_{\mathbb{R}}\left|3\sqrt{6}\gamma^2P_{\varepsilon,\beta}^2u_{\varepsilon,\beta}\right|\left|\frac{u_{\varepsilon,\beta}^3}{\sqrt{6}}\right|dx+\frac{1}{12}\left\|u_{\varepsilon,\beta}(t,\cdot)\right\|_{L^6(\mathbb{R})}^6\\
&\le27\gamma^4\int_{\mathbb{R}}P_{\varepsilon,\beta}^4u_{\varepsilon,\beta}^2dx+\frac{1}{6}\left\|u_{\varepsilon,\beta}(t,\cdot)\right\|_{L^6(\mathbb{R})}^6\\
&\le27\gamma^4\left\|P_{\varepsilon,\beta}\right\|_{L^\infty((0,T)\times\mathbb{R})}^4\left\|u_{\varepsilon,\beta}(t,\cdot)\right\|_{L^2(\mathbb{R})}^2+\frac{1}{6}\left\|u_{\varepsilon,\beta}(t,\cdot)\right\|_{L^6(\mathbb{R})}^6\\
&\le27\gamma^4C_0e^{2\gamma t}\left\|P_{\varepsilon,\beta}\right\|_{L^\infty((0,T)\times\mathbb{R})}^4+\frac{1}{6}\left\|u_{\varepsilon,\beta}(t,\cdot)\right\|_{L^6(\mathbb{R})}^6\\
&\le C(T)\left\|P_{\varepsilon,\beta}\right\|_{L^\infty((0,T)\times\mathbb{R})}^4+\frac{1}{6}\left\|u_{\varepsilon,\beta}(t,\cdot)\right\|_{L^6(\mathbb{R})}^6.
\end{aligned}\tag{2.50}$$

Since $0 < \varepsilon < 1$, it follows from (2.1), (2.7) and the Young inequality that

$$\begin{aligned}
&-3\gamma\varepsilon^2\int_{\mathbb{R}}\partial_{xx}^2u_{\varepsilon,\beta}P_{\varepsilon,\beta}dx=3\gamma\varepsilon^2\int_{\mathbb{R}}\partial_xu_{\varepsilon,\beta}\partial_xP_{\varepsilon,\beta}dx\\
&\leq 3\gamma\varepsilon^2\left|\int_{\mathbb{R}}\partial_xu_{\varepsilon,\beta}\partial_xP_{\varepsilon,\beta}dx\right|\leq 3\varepsilon^2\int_{\mathbb{R}}\left|\partial_xu_{\varepsilon,\beta}\right|\left|\gamma\partial_xP_{\varepsilon,\beta}\right|dx\\
&\leq\frac{3\varepsilon^2}{2}\left\|\partial_xu_{\varepsilon,\beta}(t,\cdot)\right\|_{L^2(\mathbb{R})}^2+\frac{3\gamma^2\varepsilon^2}{2}\left\|\partial_xP_{\varepsilon,\beta}(t,\cdot)\right\|_{L^2(\mathbb{R})}^2\\
&\leq\frac{3\varepsilon^2}{2}\left\|\partial_xu_{\varepsilon,\beta}(t,\cdot)\right\|_{L^2(\mathbb{R})}^2+\frac{3\gamma^2\varepsilon^2}{2}C_0e^{2\gamma t}\\
&\leq\frac{3\varepsilon^2}{2}\left\|\partial_xu_{\varepsilon,\beta}(t,\cdot)\right\|_{L^2(\mathbb{R})}^2+\frac{3\gamma^2}{2}C_0e^{2\gamma t}\\
&\leq\frac{3\varepsilon}{2}\left\|\partial_xu_{\varepsilon,\beta}(t,\cdot)\right\|_{L^2(\mathbb{R})}^2+C(T).
\end{aligned}\tag{2.51}$$

Thanks to (1.13), (2.16) and the Young inequality,

$$\begin{aligned}
&10\beta\left|\int_{\mathbb{R}}u_{\varepsilon,\beta}^3(\partial_xu_{\varepsilon,\beta})^2dx\right|\leq 10\beta\int_{\mathbb{R}}|u_{\varepsilon,\beta}|^3(\partial_xu_{\varepsilon,\beta})^2dx\\
&\leq 10\beta\left\|u_{\varepsilon,\beta}\right\|_{L^\infty((0,T)\times\mathbb{R})}\int_{\mathbb{R}}u_{\varepsilon,\beta}^2(\partial_xu_{\varepsilon,\beta})^2dx\leq C(T)\beta^{\frac{1}{2}}\int_{\mathbb{R}}u_{\varepsilon,\beta}^2(\partial_xu_{\varepsilon,\beta})^2dx\\
&\leq\varepsilon\int_{\mathbb{R}}|C_0C(T)\partial_xu_{\varepsilon,\beta}|u_{\varepsilon,\beta}^2|\partial_xu_{\varepsilon,\beta}|dx\\
&\leq\varepsilon C(T)\left\|\partial_xu_{\varepsilon,\beta}(t,\cdot)\right\|_{L^2(\mathbb{R})}^2+\frac{\varepsilon}{2}\int_{\mathbb{R}}u_{\varepsilon,\beta}^4(\partial_xu_{\varepsilon,\beta})^2dx.
\end{aligned}\tag{2.52}$$

For the Young inequality,

$$\begin{aligned}
&\frac{3\varepsilon^2}{2}\left|\int_{\mathbb{R}}u_{\varepsilon,\beta}^2\partial_xu_{\varepsilon,\beta}\partial_{xx}^2u_{\varepsilon,\beta}dx\right|\leq\frac{3}{2}\int_{\mathbb{R}}\left|\varepsilon^{\frac{1}{2}}u_{\varepsilon,\beta}^2\partial_xu_{\varepsilon,\beta}\right|\left|\varepsilon^{\frac{3}{2}}\partial_{xx}^2u_{\varepsilon,\beta}\right|dx\\
&\leq\frac{3\varepsilon}{4}\int_{\mathbb{R}}u_{\varepsilon,\beta}^4(\partial_xu_{\varepsilon,\beta})^2dx+\frac{3\varepsilon^3}{4}\left\|\partial_{xx}^2u_{\varepsilon,\beta}(t,\cdot)\right\|_{L^2(\mathbb{R})}^2.
\end{aligned}\tag{2.53}$$

It follows from (2.49), (2.50), (2.51) and (2.53) that

$$\begin{aligned}
&\frac{d}{dt}G_2(t)+\frac{15\varepsilon}{4}\int_{\mathbb{R}}u_{\varepsilon,\beta}^4(\partial_xu_{\varepsilon,\beta})^2dx+\frac{9\varepsilon^3}{4}\left\|\partial_{xx}^2u_{\varepsilon,\beta}(t,\cdot)\right\|_{L^2(\mathbb{R})}^2\\
&\leq G_2(t)+C(T)+C(T)\left\|P_{\varepsilon,\beta}\right\|_{L^\infty((0,T)\times\mathbb{R})}^4+\varepsilon C(T)\left\|\partial_xu_{\varepsilon,\beta}(t,\cdot)\right\|_{L^2(\mathbb{R})}^2,
\end{aligned}$$

that is

$$\begin{aligned}
&\frac{d}{dt}G_2(t)-G_2(t)+\frac{15\varepsilon}{4}\int_{\mathbb{R}}u_{\varepsilon,\beta}^4(\partial_xu_{\varepsilon,\beta})^2dx+\frac{9\varepsilon^3}{4}\left\|\partial_{xx}^2u_{\varepsilon,\beta}(t,\cdot)\right\|_{L^2(\mathbb{R})}^2\\
&\leq C(T)+C(T)\left\|P_{\varepsilon,\beta}\right\|_{L^\infty((0,T)\times\mathbb{R})}^4+\varepsilon C(T)\left\|\partial_xu_{\varepsilon,\beta}(t,\cdot)\right\|_{L^2(\mathbb{R})}^2,
\end{aligned}\tag{2.54}$$

where

$$G_2(t)=\frac{1}{6}\left\|u_{\varepsilon,\beta}(t,\cdot)\right\|_{L^6(\mathbb{R})}^6+\frac{3\varepsilon^2}{2}\left\|\partial_xu_{\varepsilon,\beta}(t,\cdot)\right\|_{L^2(\mathbb{R})}^2.\tag{2.55}$$

The Gronwall Lemma and (1.12) give

$$\begin{aligned}
&G_2(t)+\frac{15\varepsilon}{4}e^t\int_0^t\int_{\mathbb{R}} e^{-s}u_{\varepsilon,\beta}^4(s,\cdot)(\partial_x u_{\varepsilon,\beta}(s,\cdot))^2dsdx\\
&\quad+\frac{9\varepsilon^3}{4}e^t\int_0^t e^{-s}\left\|\partial_{xx}^2u_{\varepsilon,\beta}(s,\cdot)\right\|_{L^2(\mathbb{R})}^2ds\\
\leq& C_0e^t+C(T)e^t\int_0^t e^{-s}ds+C(T)e^t\left\|P_{\varepsilon,\beta}\right\|_{L^\infty((0,T)\times\mathbb{R})}^4\int_0^t e^{-s}ds\\
&\quad+\varepsilon C(T)e^t\int_0^t e^{-s}\left\|\partial_x u_{\varepsilon,\beta}(s,\cdot)\right\|_{L^2(\mathbb{R})}^2ds\\
\leq& C(T)+C(T)\left\|P_{\varepsilon,\beta}\right\|_{L^\infty((0,T)\times\mathbb{R})}^4+\varepsilon C(T)\int_0^t\left\|\partial_x u_{\varepsilon,\beta}(s,\cdot)\right\|_{L^2(\mathbb{R})}^2ds.
\end{aligned}\tag{2.56}$$

Due to (2.6),

$$\begin{aligned}
\varepsilon\int_0^t\left\|\partial_x u_{\varepsilon,\beta}(s,\cdot)\right\|_{L^2(\mathbb{R})}^2ds\leq&\varepsilon e^{2\gamma t}\int_0^t e^{-2\gamma s}\left\|\partial_x u_{\varepsilon,\beta}(s,\cdot)\right\|_{L^2(\mathbb{R})}^2ds\\
\leq&C_0e^{2\gamma t}\leq C(T).
\end{aligned}\tag{2.57}$$

Therefore, for (2.55), (2.56) and (2.57) we have

$$\begin{aligned}
&\frac{1}{6}\left\|u_{\varepsilon,\beta}(t,\cdot)\right\|_{L^6(\mathbb{R})}^6+\frac{3\varepsilon^2}{2}\left\|\partial_x u_{\varepsilon,\beta}(t,\cdot)\right\|_{L^2(\mathbb{R})}^2\\
&\qquad+\frac{15\varepsilon}{4}e^t\int_0^t\int_{\mathbb{R}} e^{-s}u_{\varepsilon,\beta}^4(s,\cdot)(\partial_x u_{\varepsilon,\beta}(s,\cdot))^2dsdx\\
&\qquad+\frac{9\varepsilon^3}{4}e^t\int_0^t e^{-s}\left\|\partial_{xx}^2u_{\varepsilon,\beta}(s,\cdot)\right\|_{L^2(\mathbb{R})}^2ds\\
\leq&C(T)+C(T)\left\|P_{\varepsilon,\beta}\right\|_{L^\infty((0,T)\times\mathbb{R})}^4.
\end{aligned}\tag{2.58}$$

Dividing (2.24) by γ, we get

$$\begin{aligned}
&\frac{d}{dt}\left(\left\|P_{\varepsilon,\beta}(t,\cdot)\right\|_{L^2(\mathbb{R})}^2+\varepsilon^2\left\|\partial_x P_{\varepsilon,\beta}(t,\cdot)\right\|_{L^2(\mathbb{R})}^2\right)\\
&\quad=-2\varepsilon\gamma\int_{\mathbb{R}}F_{\varepsilon,\beta}\partial_x P_{\varepsilon,\beta}dx+\frac{1}{3}\int_{\mathbb{R}}u_{\varepsilon,\beta}^3P_{\varepsilon,\beta}dx\\
&\qquad-\frac{\varepsilon}{3}\int_{\mathbb{R}}u_{\varepsilon,\beta}^3\partial_x P_{\varepsilon,\beta}dx+2\beta\int_{\mathbb{R}}\partial_{xx}^2u_{\varepsilon,\beta}P_{\varepsilon,\beta}dx\\
&\qquad-2\beta\varepsilon\int_{\mathbb{R}}\partial_{xx}^2u_{\varepsilon,\beta}\partial_x P_{\varepsilon,\beta}dx+2\varepsilon\int_{\mathbb{R}}\partial_x u_{\varepsilon,\beta}P_{\varepsilon,\beta}dx\\
&\qquad-2\varepsilon^2\int_{\mathbb{R}}\partial_x u_{\varepsilon,\beta}\partial_x P_{\varepsilon,\beta}dx.
\end{aligned}\tag{2.59}$$

Since $0<\varepsilon<1$, due to (2.1) and (2.11),

$$-2\varepsilon\gamma\int_{\mathbb{R}}F_{\varepsilon,\beta}\partial_x P_{\varepsilon,\beta}dx=2\varepsilon\gamma\int_{\mathbb{R}}\partial_x F_\varepsilon P_\varepsilon dx=2\varepsilon\gamma\int_{\mathbb{R}}P_{\varepsilon,\beta}^2dx\leq 2\gamma\left\|P_{\varepsilon,\beta}(t,\cdot)\right\|_{L^2(\mathbb{R})}^2.\tag{2.60}$$

It follows from (1.13), (2.1) and the Young inequality that

$$
\begin{aligned}
2\beta \int_{\mathbb{R}} \partial_{xx}^2 u_{\varepsilon,\beta} P_{\varepsilon,\beta} dx =& -2\beta \int_{\mathbb{R}} \partial_x u_{\varepsilon,\beta} \partial_x P_{\varepsilon,\beta} dx \\
\leq& 2\beta \left| \int_{\mathbb{R}} \partial_x u_{\varepsilon,\beta} \partial_x P_{\varepsilon,\beta} dx \right| \\
\leq& \varepsilon^2 \int_{\mathbb{R}} \left| \frac{C_0 \partial_x u_{\varepsilon,\beta}}{\sqrt{\gamma}} \right| |2\sqrt{\gamma} \partial_x P_{\varepsilon,\beta}| \, dx \\
\leq& \frac{C_0 \varepsilon^2}{2\gamma} \left\| \partial_x u_{\varepsilon,\beta}(t,\cdot) \right\|_{L^2(\mathbb{R})}^2 + 2\gamma \varepsilon^2 \left\| \partial_x P_{\varepsilon,\beta}(t,\cdot) \right\|_{L^2(\mathbb{R})}^2 \\
\leq& \frac{C_0 \varepsilon}{2\gamma} \left\| \partial_x u_{\varepsilon,\beta}(t,\cdot) \right\|_{L^2(\mathbb{R})}^2 + 2\gamma \varepsilon^2 \left\| \partial_x P_{\varepsilon,\beta}(t,\cdot) \right\|_{L^2(\mathbb{R})}^2 .
\end{aligned} \tag{2.61}
$$

Thanks to (1.13), (2.1), (2.7) and the Young inequality,

$$
\begin{aligned}
-2\beta\varepsilon \int_{\mathbb{R}} \partial_{xx}^2 u_{\varepsilon,\beta} \partial_x P_{\varepsilon,\beta} dx =& 2\beta\varepsilon \int_{\mathbb{R}} \partial_x u_{\varepsilon,\beta} \partial_{xx}^2 P_{\varepsilon,\beta} dx \\
\leq& 2\beta\varepsilon \left| \int_{\mathbb{R}} \partial_x u_{\varepsilon,\beta} \partial_{xx}^2 P_{\varepsilon,\beta} dx \right| \\
\leq& 2\varepsilon^3 \int_{\mathbb{R}} |C_0 \partial_x u_{\varepsilon,\beta}| \, |\partial_{xx}^2 P_{\varepsilon,\beta}| dx \\
\leq& \varepsilon^3 C_0 \left\| \partial_x u_{\varepsilon,\beta}(t,\cdot) \right\|_{L^2(\mathbb{R})}^2 + \varepsilon^3 \left\| \partial_{xx}^2 P_{\varepsilon,\beta}(t,\cdot) \right\|_{L^2(\mathbb{R})}^2 \\
\leq& \varepsilon C_0 \left\| \partial_x u_{\varepsilon,\beta}(t,\cdot) \right\|_{L^2(\mathbb{R})}^2 + C_0 e^{2\gamma t} \\
\leq& \varepsilon C_0 \left\| \partial_x u_{\varepsilon,\beta}(t,\cdot) \right\|_{L^2(\mathbb{R})}^2 + C(T).
\end{aligned} \tag{2.62}
$$

It follows from (2.1), (2.6), (2.7) and the Young inequality that

$$
\begin{aligned}
& 2\varepsilon \int_{\mathbb{R}} \partial_x u_{\varepsilon,\beta} P_{\varepsilon,\beta} dx - 2\varepsilon^2 \int_{\mathbb{R}} \partial_x u_{\varepsilon,\beta} \partial_x P_{\varepsilon,\beta} dx \\
&= -2\varepsilon \int_{\mathbb{R}} u_{\varepsilon,\beta} \partial_x P_{\varepsilon,\beta} dx + 2\varepsilon^2 \int_{\mathbb{R}} u_{\varepsilon,\beta} \partial_{xx}^2 P_{\varepsilon,\beta} dx \\
&\leq \left| -2\varepsilon \int_{\mathbb{R}} u_{\varepsilon,\beta} \partial_x P_{\varepsilon,\beta} dx + 2\varepsilon^2 \int_{\mathbb{R}} u_{\varepsilon,\beta} \partial_{xx}^2 P_{\varepsilon,\beta} dx \right| \\
&\leq 2\varepsilon \int_{\mathbb{R}} |u_{\varepsilon,\beta}| |\partial_x P_{\varepsilon,\beta}| dx + 2\varepsilon^2 \int_{\mathbb{R}} |u_{\varepsilon,\beta}| |\partial_{xx}^2 P_{\varepsilon,\beta}| dx \\
&\leq 2 \left\| u_{\varepsilon,\beta}(t,\cdot) \right\|_{L^2(\mathbb{R})}^2 + \left\| \partial_x P_{\varepsilon,\beta}(t,\cdot) \right\|_{L^2(\mathbb{R})}^2 + \varepsilon \left\| \partial_{xx}^2 P_{\varepsilon,\beta}(t,\cdot) \right\|_{L^2(\mathbb{R})}^2 \\
&\leq 3C_0 e^{2\gamma t} \leq C(T).
\end{aligned} \tag{2.63}
$$

Therefore, (2.59), (2.60), (2.61), (2.62) and (2.63) give

$$
\begin{aligned}
& \frac{d}{dt} \left(\left\| P_{\varepsilon,\beta}(t,\cdot) \right\|_{L^2(\mathbb{R})}^2 + \varepsilon^2 \left\| \partial_x P_{\varepsilon,\beta}(t,\cdot) \right\|_{L^2(\mathbb{R})}^2 \right) \\
&\qquad - 2\gamma \left(\left\| P_{\varepsilon,\beta}(t,\cdot) \right\|_{L^2(\mathbb{R})}^2 + \varepsilon^2 \left\| \partial_x P_{\varepsilon,\beta}(t,\cdot) \right\|_{L^2(\mathbb{R})}^2 \right) \\
&\leq C(T) + \left(\frac{1}{\gamma} + 1 \right) C_0 \varepsilon \left\| \partial_x u_{\varepsilon,\beta}(t,\cdot) \right\|_{L^2(\mathbb{R})}^2 \\
&\qquad + \frac{1}{3} \int_{\mathbb{R}} |u_{\varepsilon,\beta}|^3 |P_{\varepsilon,\beta}| dx + \frac{\varepsilon}{3} \int_{\mathbb{R}} |u_{\varepsilon,\beta}|^3 |\partial_x P_{\varepsilon,\beta}| dx.
\end{aligned} \tag{2.64}
$$

Due to (2.6), (2.7) (2.58) and the Young inequality,

$$
\begin{aligned}
&\frac{1}{3}\int_{\mathbb{R}}|u_{\varepsilon,\beta}|^3|P_{\varepsilon,\beta}|dx+\frac{\varepsilon}{3}\int_{\mathbb{R}}|u_{\varepsilon,\beta}|^3|\partial_x P_{\varepsilon,\beta}|dx\\
&\quad\le\frac{1}{6}\int_{\mathbb{R}}u_{\varepsilon,\beta}^2P_{\varepsilon,\beta}^2dx+\frac{1}{6}\int_{\mathbb{R}}u_{\varepsilon,\beta}^4dx\\
&\qquad+\frac{\varepsilon^2}{6}\int_{\mathbb{R}}u_{\varepsilon,\beta}^2(\partial_xP_{\varepsilon,\beta})^2dx+\frac{1}{6}\int_{\mathbb{R}}u_{\varepsilon,\beta}^4dx\\
&\quad\le\frac{1}{6}\left\|P_{\varepsilon,\beta}\right\|^2_{L^\infty((0,T)\times\mathbb{R})}\left\|u_{\varepsilon,\beta}(t,\cdot)\right\|^2_{L^2(\mathbb{R})}+\frac{\varepsilon^2}{6}\left\|\partial_xP_{\varepsilon,\beta}(t,\cdot)\right\|^2_{L^\infty(\mathbb{R})}\left\|u_{\varepsilon,\beta}(t,\cdot)\right\|^2_{L^2(\mathbb{R})}\\
&\qquad+\frac{1}{3}\int_{\mathbb{R}}\left|\frac{u_{\varepsilon,\beta}}{\sqrt{A(T)}}\right|\left|\sqrt{A(T)}u_{\varepsilon,\beta}^3\right|dx\\
&\quad\le\frac{C_0e^{2\gamma t}}{6}\left\|P_{\varepsilon,\beta}\right\|^2_{L^\infty((0,T)\times\mathbb{R})}+\frac{C_0e^{4\gamma t}}{6}+\frac{1}{6A(T)}\left\|u_{\varepsilon,\beta}(t,\cdot)\right\|^2_{L^2(\mathbb{R})}\\
&\qquad+\frac{A(T)}{6}\left\|u_{\varepsilon,\beta}(t,\cdot)\right\|^6_{L^6(\mathbb{R})}\\
&\quad\le C(T)\left\|P_{\varepsilon,\beta}\right\|^2_{L^\infty((0,T)\times\mathbb{R})}+C(T)+\frac{C_0e^{2\gamma t}}{6A(T)}+C(T)A(T)\\
&\qquad+C(T)A(T)\left\|P_{\varepsilon,\beta}\right\|^4_{L^\infty((0,T)\times\mathbb{R})}\\
&\quad\le C(T)+\frac{C(T)}{A(T)}+C(T)\left\|P_{\varepsilon,\beta}\right\|^2_{L^\infty((0,T)\times\mathbb{R})}+C(T)A(T)\left\|P_{\varepsilon,\beta}\right\|^4_{L^\infty((0,T)\times\mathbb{R})},
\end{aligned}
\tag{2.65}
$$

where $A(T)$ is a positive constant which will be specified later.

Thus, (2.64) and (2.65) give

$$
\begin{aligned}
&\frac{d}{dt}\left(\left\|P_{\varepsilon,\beta}(t,\cdot)\right\|^2_{L^2(\mathbb{R})}+\varepsilon^2\left\|\partial_xP_{\varepsilon,\beta}(t,\cdot)\right\|^2_{L^2(\mathbb{R})}\right)\\
&\qquad-2\gamma\left(\left\|P_{\varepsilon,\beta}(t,\cdot)\right\|^2_{L^2(\mathbb{R})}+\varepsilon^2\left\|\partial_xP_{\varepsilon,\beta}(t,\cdot)\right\|^2_{L^2(\mathbb{R})}\right)\\
&\quad\le C(T)+\frac{C(T)}{A(T)}+\left(\frac{1}{\gamma}+1\right)C_0\varepsilon\left\|\partial_xu_{\varepsilon,\beta}(t,\cdot)\right\|^2_{L^2(\mathbb{R})}\\
&\qquad+C(T)\left\|P_{\varepsilon,\beta}\right\|^2_{L^\infty((0,T)\times\mathbb{R})}+C(T)A(T)\left\|P_{\varepsilon,\beta}\right\|^4_{L^\infty((0,T)\times\mathbb{R})}.
\end{aligned}
$$

It follows from (1.12), (2.6) and the Gronwall Lemma that

$$
\begin{aligned}
&\left\|P_{\varepsilon,\beta}(t,\cdot)\right\|^2_{L^2(\mathbb{R})}+\varepsilon^2\left\|\partial_xP_{\varepsilon,\beta}(t,\cdot)\right\|^2_{L^2(\mathbb{R})}\\
&\quad\le C_0e^{2\gamma t}+C(T)e^{2\gamma t}\int_0^t e^{-2\gamma s}ds+\frac{C(T)}{A(T)}e^{2\gamma t}\int_0^t e^{-2\gamma s}ds\\
&\qquad+\left(\frac{1}{\gamma}+1\right)C_0e^{2\gamma t}\varepsilon\int_0^t e^{-2\gamma s}\left\|\partial_xu_{\varepsilon,\beta}(s,\cdot)\right\|^2_{L^2(\mathbb{R})}\\
&\qquad+C(T)e^{2\gamma t}\left\|P_{\varepsilon,\beta}\right\|^2_{L^\infty((0,T)\times\mathbb{R})}\int_0^t e^{-2\gamma s}ds\\
&\qquad+C(T)A(T)e^{2\gamma t}\left\|P_{\varepsilon,\beta}\right\|^4_{L^\infty((0,T)\times\mathbb{R})}\int_0^t e^{-2\gamma s}ds\\
&\quad\le C(T)+\frac{C(T)}{A(T)}+\left(\frac{1}{\gamma}+1\right)\frac{C_0e^{2\gamma t}}{2}+C(T)\left\|P_{\varepsilon,\beta}\right\|^2_{L^\infty((0,T)\times\mathbb{R})}
\end{aligned}
$$

$$+ C(T)A(T) \left\| P_{\varepsilon,\beta} \right\|^4_{L^\infty((0,T)\times\mathbb{R})},$$

that is

$$\begin{aligned} &\left\| P_{\varepsilon,\beta}(t,\cdot) \right\|^2_{L^2(\mathbb{R})} + \varepsilon^2 \left\| \partial_x P_{\varepsilon,\beta}(t,\cdot) \right\|^2_{L^2(\mathbb{R})} \\ &\quad \leq C(T) \left(1 + \frac{1}{A(T)} + \left\| P_{\varepsilon,\beta} \right\|^2_{L^\infty((0,T)\times\mathbb{R})} + A(T) \left\| P_{\varepsilon,\beta} \right\|^4_{L^\infty((0,T)\times\mathbb{R})} \right). \end{aligned} \tag{2.66}$$

We prove (2.39). Due to the Hölder inequality,

$$P^2_{\varepsilon,\beta}(t,x) \leq 2 \int_{\mathbb{R}} |P_{\varepsilon,\beta} \partial_x P_{\varepsilon,\beta}| dx \leq \left\| P_{\varepsilon,\beta}(t,\cdot) \right\|_{L^2(\mathbb{R})} \left\| \partial_x P_{\varepsilon,\beta}(t,\cdot) \right\|_{L^2(\mathbb{R})},$$

that is

$$P^4_{\varepsilon,\beta}(t,x) \leq 4 \left\| P_{\varepsilon,\beta}(t,\cdot) \right\|^2_{L^2(\mathbb{R})} \left\| \partial_x P_{\varepsilon,\beta}(t,\cdot) \right\|^2_{L^2(\mathbb{R})}.$$

For (2.7) and (2.66),

$$P^4_{\varepsilon,\beta}(t,x) \leq C(T) \left(1 + \frac{1}{A(T)} + \left\| P_{\varepsilon,\beta} \right\|^2_{L^\infty((0,T)\times\mathbb{R})} + A(T) \left\| P_{\varepsilon,\beta} \right\|^4_{L^\infty((0,T)\times\mathbb{R})} \right).$$

Therefore,

$$(1 - C(T)A(T)) \left\| P_{\varepsilon,\beta} \right\|^4_{L^\infty((0,T)\times\mathbb{R})} - C(T) \left\| P_{\varepsilon,\beta} \right\|^2_{L^\infty((0,T)\times\mathbb{R})} - C(T) - \frac{C(T)}{A(T)} \leq 0.$$

Choosing

$$A(T) = \frac{1}{2C(T)}, \tag{2.67}$$

we have that

$$\frac{1}{2} \left\| P_{\varepsilon,\beta} \right\|^4_{L^\infty((0,T)\times\mathbb{R})} - C(T) \left\| P_{\varepsilon,\beta} \right\|^2_{L^\infty((0,T)\times\mathbb{R})} - C(T) \leq 0,$$

which gives (2.39). Therefore, (2.39) and (2.67) give (2.40) and (2.41), while (2.42), (2.43), (2.44) and (2.45) follow from (2.39).

We show that (2.47) holds true. We begin by observing that, for (2.45),

$$\varepsilon^3 \int_0^t \left\| \partial^2_{xx} u_{\varepsilon,\beta}(s,\cdot) \right\|^2_{L^2(\mathbb{R})} ds \leq \varepsilon^3 e^t \int_0^t e^{-s} \left\| \partial^2_{xx} u_{\varepsilon,\beta}(s,\cdot) \right\|^2_{L^2(\mathbb{R})} ds \leq C(T).$$

Therefore,

$$\varepsilon^3 \int_0^T \left\| \partial^2_{xx} u_{\varepsilon,\beta}(t,\cdot) \right\|^2_{L^2(\mathbb{R})} dt \leq C(T). \tag{2.68}$$

Moreover, for (2.57),

$$\varepsilon \int_0^T \left\| \partial_x u_{\varepsilon,\beta}(t,\cdot) \right\|^2_{L^2(\mathbb{R})} dt \leq C(T). \tag{2.69}$$

Thus, thanks to (1.13), (2.68), (2.69) and Hölder inequality,

$$\begin{aligned} \beta \int_0^T \int_{\mathbb{R}} |\partial_x u_{\varepsilon,\beta} \partial^2_{xx} u_{\varepsilon,\beta}| ds dx &= \frac{\beta}{\varepsilon^2} \int_0^T \int_{\mathbb{R}} \varepsilon^{\frac{1}{2}} |\partial_x u_{\varepsilon,\beta}| \varepsilon^{\frac{3}{2}} |\partial^2_{xx} u_{\varepsilon,\beta}| dx \\ &\leq \frac{\beta}{\varepsilon^2} \left(\varepsilon \int_0^T \int_{\mathbb{R}} (\partial_x u_{\varepsilon,\beta})^2 ds dx \right)^{\frac{1}{2}} \left(\varepsilon^3 \int_0^T \int_{\mathbb{R}} (\partial^2_{xx} u_{\varepsilon,\beta})^2 ds dx \right)^{\frac{1}{2}} \\ &\leq C(T) \frac{\beta}{\varepsilon^2} \leq C(T), \end{aligned}$$

that is (2.46).

Finally, we prove (2.47). Due to (1.13) and (2.68), we have

$$\beta^2 \int_0^T \left\|\partial_{xx}^2 u_{\varepsilon,\beta}(s,\cdot)\right\|_{L^2(\mathbb{R})}^2 ds \le C_0^2 \varepsilon^4 \int_0^T \left\|\partial_{xx}^2 u_{\varepsilon,\beta}(s,\cdot)\right\|_{L^2(\mathbb{R})}^2 ds \le C(T)\varepsilon,$$

which gives (2.47). □

3. Proof of Theorem 1.1

In this section, we prove Theorem 1.1. The following technical lemma is needed [14].

Lemma 3.1. *Let Ω be a bounded open subset of $\mathbb{R}^2$. Suppose that the sequence $\{\mathcal{L}_n\}_{n\in\mathbb{N}}$ of distributions is bounded in $W^{-1,\infty}(\Omega)$. Suppose also that*

$$\mathcal{L}_n = \mathcal{L}_{1,n} + \mathcal{L}_{2,n},$$

where $\{\mathcal{L}_{1,n}\}_{n\in\mathbb{N}}$ lies in a compact subset of $H^{-1}_{loc}(\Omega)$ and $\{\mathcal{L}_{2,n}\}_{n\in\mathbb{N}}$ lies in a bounded subset of $\mathcal{M}_{loc}(\Omega)$. Then $\{\mathcal{L}_n\}_{n\in\mathbb{N}}$ lies in a compact subset of $H^{-1}_{loc}(\Omega)$.

Moreover, we consider the following definition.

Definition 3.1. *A pair of functions (η, q) is called an entropy–entropy flux pair if $\eta : \mathbb{R} \to \mathbb{R}$ is a C^2 function and $q : \mathbb{R} \to \mathbb{R}$ is defined by*

$$q(u) = -\int_0^u \frac{\xi^2}{2}\eta'(\xi)d\xi.$$

An entropy-entropy flux pair (η, q) is called convex/compactly supported if, in addition, η is convex/compactly supported.

We begin by proving the following result.

Lemma 3.2. *Assume that (1.7), (1.8), (1.9), (1.12), and (1.13) hold. Then for any compactly supported entropy–entropy flux pair (η, q), there exist two sequences $\{\varepsilon_n\}_{n\in\mathbb{N}}$, $\{\beta_n\}_{n\in\mathbb{N}}$, with $\varepsilon_n, \beta_n \to 0$, and two limit functions*

$$u \in L^\infty(0,T; L^2(\mathbb{R}) \cap L^6(\mathbb{R})),$$
$$P \in L^\infty((0,T)\times\mathbb{R}) \cap L^2((0,T)\times\mathbb{R}),$$

such that

$$u_{\varepsilon_n,\beta_n} \to u \quad \text{in} \quad L^p_{loc}((0,\infty)\times\mathbb{R}), \quad \text{for each} \quad 1 \le p < 6, \tag{3.1}$$

$$P_{\varepsilon_n,\beta_n} \to P \quad \text{in} \quad L^\infty((0,T)\times\mathbb{R}) \cap L^2((0,T)\times\mathbb{R}), \tag{3.2}$$

and u is a distributional solution of (1.4).

Proof. Let us consider a compactly supported entropy–entropy flux pair (η, q). Multiplying (1.11) by $\eta'(u_{\varepsilon,\beta})$, we have

$$\begin{aligned}\partial_t \eta(u_{\varepsilon,\beta}) + \partial_x q(u_{\varepsilon,\beta}) =& \varepsilon\eta'(u_{\varepsilon,\beta})\partial_{xx}^2 u_{\varepsilon,\beta} + \beta\eta'(u_{\varepsilon,\beta})\partial_{xxx}^3 u_{\varepsilon,\beta} + \gamma\eta'(u_{\varepsilon,\beta})P_{\varepsilon,\beta}\\ =& I_{1,\varepsilon,\beta} + I_{2,\varepsilon,\beta} + I_{3,\varepsilon,\beta} + I_{4,\varepsilon,\beta} + I_{5,\varepsilon,\beta},\end{aligned}$$

where

$$\begin{aligned} I_{1,\varepsilon,\beta} &= \partial_x(\varepsilon\eta'(u_{\varepsilon,\beta})\partial_x u_{\varepsilon,\beta}),\\ I_{2,\varepsilon,\beta} &= -\varepsilon\eta''(u_{\varepsilon,\beta})(\partial_x u_{\varepsilon,\beta})^2,\\ I_{3,\varepsilon,\beta} &= \partial_x(\beta\eta'(u_{\varepsilon,\beta})\partial_{xx}^2 u_{\varepsilon,\beta}),\\ I_{4,\varepsilon,\beta} &= -\beta\eta''(u_{\varepsilon,\beta})\partial_x u_{\varepsilon,\beta}\partial_{xx}^2 u_{\varepsilon,\beta},\\ I_{5,\varepsilon,\beta} &= \gamma\eta'(u_{\varepsilon,\beta})P_{\varepsilon,\beta}. \end{aligned} \tag{3.3}$$

Arguing as [6, Lemma 3.2], we have that $I_{1,\varepsilon,\beta} \to 0$ in $H^{-1}((0,T)\times\mathbb{R})$, $\{I_{2,\varepsilon,\beta}\}_{\varepsilon,\beta>0}$ is bounded in $L^1((0,T)\times\mathbb{R})$, $I_{3,\varepsilon,\beta} \to 0$ in $H^{-1}((0,T)\times\mathbb{R})$, $\{I_{4,\varepsilon,\beta}\}_{\varepsilon,\beta>0}$ is bounded in $L^1((0,T)\times\mathbb{R})$, and $\{I_{5,\varepsilon,\beta}\}_{\varepsilon,\beta>0}$ is bounded in $L^1_{loc}((0,\infty)\times\mathbb{R})$. Therefore, Lemma 3.1 and the L^p compensated compactness of [17] give (3.1). (3.2) follows from Lemma 2.4.

We conclude by proving that u is a distributional solution of (1.4). Let $\phi \in C^\infty(\mathbb{R}^2)$ be a test function with compact support. We have to prove that

$$\int_0^\infty\int_{\mathbb{R}} \left(u\partial_t\phi - \frac{u^3}{6}\partial_x\phi \right) dtdx - \gamma \int_0^\infty\int_{\mathbb{R}} P\phi dtdx + \int_{\mathbb{R}} u_0(x)\phi(0,x)dx = 0. \tag{3.4}$$

We have that

$$\begin{aligned}
&\int_0^\infty\int_{\mathbb{R}} \left(u_{\varepsilon_n,\beta_n}\partial_t\phi - \frac{u^3_{\varepsilon_n,\beta_n}}{6}\partial_x\phi \right) dtdx - \gamma \int_0^\infty\int_{\mathbb{R}} P_{\varepsilon_n,\beta_n}\phi dtdx + \int_{\mathbb{R}} u_{0,\varepsilon_n,\beta_n}(x)\phi(0,x)dx \\
&\quad + \varepsilon_n \int_0^\infty\int_{\mathbb{R}} u_{\varepsilon_n,\beta_n}\partial^2_{xx}\phi dtdx + \varepsilon_n \int_0^\infty u_{0,\varepsilon_n,\beta_n}(x)\partial^2_{xx}\phi(0,x)dx \\
&\quad - \beta_n \int_0^\infty\int_{\mathbb{R}} u_{\varepsilon_n,\beta_n}\partial^3_{xxx}\phi dtdx - \beta_n \int_0^\infty u_{0,\varepsilon_n,\beta_n}(x)\partial^3_{xxx}\phi(0,x)dx = 0.
\end{aligned}$$

Therefore, (3.4) follows from (3.1), and (3.2). □

Lemma 3.3. *Assume that* (1.7), (1.8), (1.9), (1.12), *and* (1.14) *hold. Then,*

$$u_{\varepsilon_n,\beta_n} \to u \quad \text{in} \quad L^p_{loc}((0,\infty)\times\mathbb{R}), \quad \text{for each} \quad 1 \le p < 6, \tag{3.5}$$

$$P_{\varepsilon_n,\beta_n} \to P \quad \text{in} \quad L^p_{loc}(0,\infty;W^{1,\infty}(\mathbb{R})\cap H^1(\mathbb{R})), \quad \text{for each} \quad 1 \le p < 6, \tag{3.6}$$

where u is the unique entropy solution of (1.4).

Proof. Let us consider a compactly supported entropy–entropy flux pair (η, q). Multiplying (1.11) by $\eta'(u_{\varepsilon,\beta})$, we obtain that

$$\begin{aligned}
\partial_t\eta(u_{\varepsilon,\beta}) + \partial_x q(u_{\varepsilon,\beta}) =& \varepsilon\eta'(u_{\varepsilon,\beta})\partial^2_{xx}u_{\varepsilon,\beta} + \beta\eta'(u_{\varepsilon,\beta})\partial^3_{xxx}u_{\varepsilon,\beta} + \gamma\eta'(u_{\varepsilon,\beta})P_{\varepsilon,\beta} \\
=& I_{1,\varepsilon,\beta} + I_{2,\varepsilon,\beta} + I_{3,\varepsilon,\beta} + I_{4,\varepsilon,\beta} + I_{5,\varepsilon,\beta},
\end{aligned}$$

where $I_{1,\varepsilon,\beta}$, $I_{2,\varepsilon,\beta}$, $I_{3,\varepsilon,\beta}$, $I_{4,\varepsilon,\beta}$ and $I_{5,\varepsilon,\beta}$ are defined in (3.3).

Arguing as [6, Lemma 3.3], we obtain that $I_{1,\varepsilon,\beta} \to 0$ in $H^{-1}((0,T)\times\mathbb{R})$, $\{I_{2,\varepsilon,\beta}\}_{\varepsilon,\beta>0}$ is bounded in $L^1((0,T)\times\mathbb{R})$, $I_{3,\varepsilon,\beta} \to 0$ in $H^{-1}((0,T)\times\mathbb{R})$, $I_{4,\varepsilon,\beta} \to 0$ in $L^1((0,T)\times\mathbb{R})$, and $\{I_{5,\varepsilon,\beta}\}_{\varepsilon,\beta>0}$ is bounded in $L^1_{loc}((0,\infty)\times\mathbb{R})$. Therefore, Lemma 3.1 gives (3.5).

We prove (3.6). We begin by observing that, integrating the second equation of (1.11) on $(0,x)$, we have

$$P_{\varepsilon,\beta}(t,x) = \int_0^x u_{\varepsilon,\beta}(t,y)dy + \varepsilon\partial_x P_{\varepsilon,\beta}(t,x) - \varepsilon\partial_x P_{\varepsilon,\beta}(t,0). \tag{3.7}$$

Let us show that

$$\varepsilon\partial_x P_\varepsilon \to 0 \text{ in } L^\infty((0,T)\times\mathbb{R}),\ T>0. \tag{3.8}$$

It follows from (2.7) that

$$\varepsilon \left\|\partial_x P_\varepsilon\right\|_{L^\infty((0,T)\times\mathbb{R})} \le \sqrt{\varepsilon}e^{\gamma T}\left\|u_0\right\|^2_{L^2(\mathbb{R})} = \sqrt{\varepsilon}C(T) \to 0,$$

that is (3.8).
We claim that

$$\varepsilon\partial_x P_\varepsilon(\cdot,0) \to 0 \text{ in } L^\infty(0,T),\ T>0. \tag{3.9}$$

Again by (2.7), we have that

$$\varepsilon \left\|\partial_x P_\varepsilon(\cdot,0)\right\|_{L^\infty(0,T)} \le \sqrt{\varepsilon}e^{\gamma T}\left\|u_0\right\|^2_{L^2(\mathbb{R})} = \sqrt{\varepsilon}C(T) \to 0,$$

that is (3.9). Therefore, (3.6) follows from (3.5), (3.7), (3.8), (3.9) and the Hölder inequality.

We conclude by proving that u is the unique entropy solution of (1.4). Let us consider a compactly supported entropy–entropy flux pair (η, q), and $\phi \in C_c^\infty((0,\infty)\times\mathbb{R})$ a non–negative function. We have to prove that

$$\int_0^\infty\!\!\int_{\mathbb{R}} (\partial_t \eta(u) + \partial_x q(u))\phi dt dx - \gamma \int_0^\infty\!\!\int_{\mathbb{R}} P_{\varepsilon,\beta}\eta'(u)\phi dt dx \le 0. \tag{3.10}$$

We have that

$$\begin{aligned}
&\int_0^\infty\!\!\int_{\mathbb{R}} (\partial_x \eta(u_{\varepsilon_n,\beta_n}) + \partial_x q(u_{\varepsilon_n,\beta_n}))\phi dt dx - \gamma \int_0^\infty\!\!\int_{\mathbb{R}} P_{\varepsilon_n,\beta_n}\eta'(u_{\varepsilon_n,\beta_n})\phi dt dx \\
&\quad= \varepsilon_n \int_0^\infty\!\!\int_{\mathbb{R}} \partial_x(\eta'(u_{\varepsilon_n,\beta_n})\partial_x u_{\varepsilon_n,\beta_n})\phi dt dx - \varepsilon_n \int_0^\infty\!\!\int_{\mathbb{R}} \eta''(u_{\varepsilon_n,\beta_n})(\partial_x u_{\varepsilon_n,\beta_n})^2\phi dt dx \\
&\qquad+ \beta_n \int_0^\infty\!\!\int_{\mathbb{R}} \partial_x(\eta'(u_{\varepsilon_n,\beta_n})\partial_{xx}^2 u_{\varepsilon_n,\beta_n})\phi dt dx \\
&\qquad- \beta_n \int_0^\infty\!\!\int_{\mathbb{R}} \eta''(u_{\varepsilon_n,\beta_n})\partial_x u_{\varepsilon_n,\beta_n}\partial_{xx}^2 u_{\varepsilon_n,\beta_n}\phi dt dx \\
&\quad\le -\varepsilon_n \int_0^\infty\!\!\int_{\mathbb{R}} \eta'(u_{\varepsilon_n,\beta_n})\partial_x u_{\varepsilon_n,\beta_n}\partial_x\phi dt dx - \beta_n \int_0^\infty\!\!\int_{\mathbb{R}} \eta'(u_{\varepsilon_n,\beta_n})\partial_{xx}^2 u_{\varepsilon_n,\beta_n}\partial_x\phi dt dx \\
&\qquad- \beta_n \int_0^\infty\!\!\int_{\mathbb{R}} \eta''(u_{\varepsilon_n,\beta_n})\partial_x u_{\varepsilon_n,\beta_n}\partial_{xx}^2 u_{\varepsilon_n,\beta_n}\phi dt dx \\
&\quad\le \varepsilon_n \int_0^\infty\!\!\int_{\mathbb{R}} |\eta'(u_{\varepsilon_n,\beta_n})||\partial_x u_{\varepsilon_n,\beta_n}||\partial_x\phi| dt dx \\
&\qquad+ \beta_n \int_0^\infty\!\!\int_{\mathbb{R}} |\eta'(u_{\varepsilon_n,\beta_n})||\partial_{xx}^2 u_{\varepsilon_n,\beta_n}||\partial_x\phi| dt dx \\
&\qquad+ \beta_n \int_0^\infty\!\!\int_{\mathbb{R}} |\eta''(u_{\varepsilon_n,\beta_n})||\partial_x u_{\varepsilon_n,\beta_n}\partial_{xx}^2 u_{\varepsilon_n,\beta_n}||\phi| dt dx \\
&\quad\le \varepsilon_n \left\|\eta'\right\|_{L^\infty(\mathbb{R})} \left\|\partial_x u_{\varepsilon_n,\beta_n}\right\|_{L^2(\mathrm{supp}\,(\partial_x\phi))} \left\|\partial_x\phi\right\|_{L^2(\mathrm{supp}\,(\partial_x\phi))} \\
&\qquad+ \beta_n \left\|\eta'\right\|_{L^\infty(\mathbb{R})} \left\|\partial_{xx}^2 u_{\varepsilon_n,\beta_n}\right\|_{L^2(\mathrm{supp}\,(\partial_x\phi))} \left\|\partial_x\phi\right\|_{L^2(\mathrm{supp}\,(\partial_x\phi))} \\
&\qquad+ \beta_n \left\|\eta''\right\|_{L^\infty(\mathbb{R})} \left\|\phi\right\|_{L^\infty(\mathbb{R})} \left\|\partial_x u_{\varepsilon_n,\beta_n}\partial_{xx}^2 u_{\varepsilon_n,\beta_n}\right\|_{L^1(\mathrm{supp}\,(\partial_x\phi))} \\
&\quad\le \varepsilon_n \left\|\eta'\right\|_{L^\infty(\mathbb{R})} \left\|\partial_x u_{\varepsilon_n,\beta_n}\right\|_{L^2((0,T)\times\mathbb{R})} \left\|\partial_x\phi\right\|_{L^2((0,T)\times\mathbb{R})} \\
&\qquad+ \beta_n \left\|\eta'\right\|_{L^\infty(\mathbb{R})} \left\|\partial_{xx}^2 u_{\varepsilon_n,\beta_n}\right\|_{L^2((0,T)\times\mathbb{R})} \left\|\partial_x\phi\right\|_{L^2((0,T)\times\mathbb{R})} \\
&\qquad+ \beta_n \left\|\eta''\right\|_{L^\infty(\mathbb{R})} \left\|\phi\right\|_{L^\infty(\mathbb{R}^+\times\mathbb{R})} \left\|\partial_x u_{\varepsilon_n,\beta_n}\partial_{xx}^2 u_{\varepsilon_n,\beta_n}\right\|_{L^1((0,T)\times\mathbb{R})}.
\end{aligned}$$

(3.10) follows from (1.14), (3.5), (3.6), Lemmas 2.1 and 2.4. □

Proof of Theorem 1.1. i), *ii*), and *iii*) follow from Lemma 3.2. *iv*) and *v*) follow from Lemma 3.3, while (1.15) follows from (2.3) (3.1), or (3.5). Therefore, the proof is done. □

References

[1] J. C. Brunelli. The short pulse hierarchy. *J. Math. Phys.* 46:123507, 2005.

[2] Y. Chung, C. K. R. T. Jones, T. Schäfer, and C. E. Wayne. Ultra-short pulses in linear and nonlinear media. *Nonlinearity*, 18:1351–1374, 2005.

[3] G. M. Coclite and L. di Ruvo. Wellposedness of bounded solutions of the non-homogeneous initial boundary value problem for the Ostrovsky-Hunter equation. Submitted.

[4] G. M. Coclite and L. di Ruvo. Oleinik type estimates for the Ostrovsky–Hunter Equation. Submitted.

[5] G. M. Coclite and L. di Ruvo. Wellposedness results for the Short Pulse Equation. Submitted.

[6] G. M. Coclite and L. di Ruvo. Convergence of the Ostrovsky Equation to the Ostrovsky-Hunter One. *J. Differential Equations*, 256:3245–3277, 2014.
[7] G. M. Coclite, L. di Ruvo, and K. H. Karlsen. Some wellposedness results for the Ostrovsky-Hunter equation. To appear on *Springer Proceedings in Mathematics & Statistics*.
[8] G. M. Coclite and K. H. Karlsen. A singular limit problem for conservation laws related to the Camassa-Holm shallow water equation. *Comm. Partial Differential Equations*, 31:1253–1272, 2006.
[9] N. Costanzino, V. Manukian, and C.K.R.T. Jones Solitary waves of the regularized short pulse and Ostrovsky equations *SIAM J. Math. Anal*, 41:2088-2106, 2009.
[10] L. di Ruvo. Discontinuous solutions for the Ostrovsky–Hunter equation and two phase flows. *Phd Thesis, University of Bari*, 2013.
[11] J. Hunter and K. P. Tan. Weakly dispersive short waves *Proceedings of the IVth international Congress on Waves and Stability in Continuous Media*, Sicily, 1987.
[12] Y. Liu, D. Pelinovsky, and A. Sakovich. Wave breaking in the short-pulse equation. *Dynamics of PDE*, 6:291–310, 2009.
[13] P. G. LeFloch and R. Natalini. Conservation laws with vanishing nonlinear diffusion and dispersion. *Nonlinear Anal. 36, no. 2, Ser. A: Theory Methods*, 212–230, 1992
[14] F. Murat. L'injection du cône positif de H^{-1} dans $W^{-1,q}$ est compacte pour tout $q < 2$. *J. Math. Pures Appl. (9)*, 60(3):309–322, 1981.
[15] L. A. Ostrovsky. Nonlinear internal waves in a rotating ocean. *Okeanologia*, 18:181–191, 1978.
[16] D. Pelinovsky and A. Sakovich. Global well-posedness of the short-pulse and sine-Gordon equations in energy space. *Comm. Partial Differential Equations* 35(4), 613–629, 2010.
[17] M. E. Schonbek. Convergence of solutions to nonlinear dispersive equations *Comm. Partial Differential Equations*, 7(8):959–1000, 1982.
[18] A. Sakovich and S. Sakovich. The short pulse equation is integrable. *J. Phys. Soc. Jpn.* 74:239–241, 2005.
[19] A. Sakovich and S. Sakovich. Solitary wave solutions of the short pulse equation. *J. Phys. Soc. Jpn.* 39:361-367, 2006.
[20] T. Schäfer and C.E. Wayne. Propagation of ultra-short optical pulses in cubic nonlinear media. *Physica D*, 196:90–105, 2004.

(Giuseppe Maria Coclite and Lorenzo di Ruvo)
Department of Mathematics, University of Bari, via E. Orabona 4, 70125 Bari, Italy
E-mail address: giuseppemaria.coclite@uniba.it, lorenzo.diruvo@uniba.it
URL: http://www.dm.uniba.it/Members/coclitegm/